



# **Visual Recognition Standards Guide**

## FOR GRAIN COMMODITY SAMPLING & ASSESSMENT

Issued 1<sup>st</sup> August 2024



**Visual Recognition Standards Guide. Issued 1st August 2024**

Grain Trade Australia (GTA)

Disclaimer:

The photographs and illustrations reproduced in this Guide are produced to supplement the Grain Trade Australia Commodity Trading Standards and for the purpose of facilitating the trade of the depicted commodities, within Australia.

These images may not be used in part or full for commercial use without the expressed permission of the Grain Trade Australia Trading Standards Committee.



# TABLE OF CONTENTS

## Introduction

### Document Calibration

### Printer Calibration

## BARLEY

### Varietal Identification

#### Common Defects

Sprouted

Dark Tipped (Germ End Stained-WA)

Shot

Skinnings (Skinned-WA)

Cleaved (front, back and side)

Distorted

Dry Green or Sappy

Broken

Heavily Discoloured-WA

Pink Fungal Staining

Field Fungal (Spotted/Field Fungi Affected-WA)

Severely Damaged inc. Fusarium (except WA)

Coloured Aleurone Layer (Blue or Black)

Insect Damaged

Pickling Compounds or Artificial Colour (Pickled Barley -WA)

## CANOLA

#### Common Defects

Broken or Split

Sprouted

Heat Damaged

Weather Damaged

Mould

Frost Damaged

Green Seeds

## 1 CHICKPEAS, DESI

### 2 Common Defects

#### 3 Severely Damaged

Broken, Chipped, Loose Seed Coat and Split

Insect Damaged

Hail Damaged

#### 5 Sprouted

#### 6 Shrivelled and Wrinkled

6 Green

6 Pickling Compounds or Artificial Colouring

6 Poor Colour

7 Stained and Weather Damaged

7 Fungal Affected (e.g. Ascochyta)

8

8

## 8 CHICKPEAS, KABULI

### 9 Common Defects

9 Poor Colour

9 Insect Damaged

10 Broken, Chipped, Loose Seed Coat and Split

11 Severely Damaged

11 Sprouted

11 Shrivelled and Wrinkled

## FABA BEANS

### 13 Common Defects

13 Severely Damaged

13 Shrivelled and Wrinkled

13 Sprouted

13 Insect Damaged

14 Broken, Chipped, Loose Seed Coat and Split

14 Poor Colour

14 Green

14 Fungal Affected (e.g. Ascochyta)

16

16

17

18

18

18

19

19

19

19

21

21

21

23

23

23

23

24

25

25

25

25

27

27

28

28

28

28

29

30

30

31



## **FABA BEANS (continued...)**

Stained and Weather Damaged  
Pea Seed Borne Mosaic Virus

## **LENTILS, RED**

### **Common Defects**

Shrivelled and Wrinkled  
Broken, Chipped, Loose Seed Coat and Split  
Insect Damaged  
Sprouted  
Severely Damaged  
Poor Colour Seed Coat  
Orange Tip  
Poor Colour Kernel - Dehulled Lentil  
Blonde Kernel  
Fungal Affected  
Stained and Weather Damaged  
Contrasting Colours

## **LUPINS, ANGUSTIFOLIUS**

### **Common Defects**

Broken, Chipped, Loose Seed Coat and Split  
Manganese Deficiency  
Shrivelled and Wrinkled  
Insect Damaged  
Poor Colour (Discoloured-WA)  
Phomopsis  
Bitter Dark (Bitter Variety-WA)  
Severely Damaged  
Sprouted  
Pickling Compounds or Artificial Colour

## **MAIZE, FEED**

### **31 Common Defects**

32 Heat Damaged / Bin Burnt  
Broken  
Insect Damaged  
34 Sprouted  
34 Storage Mould  
35 Fungal Affected  
36 Dead  
36 Pickling Compounds or Artificial Colouring  
36 Field Fungi  
37 Kernel Red Streak

## **37 OATS**

### **37 Common Defects**

38 Severely Damaged  
38 Field Fungi (Spotted Mould Affected-WA)  
39 Stained Grains (Heavily Discoloured-WA)  
Septoria  
Stained Groats  
41 Insect Damaged  
41 Sprouted  
41 Damaged Grains  
42 Dry Green or Sappy  
42 Shot

## **43 PEAS, FIELD**

### **43 Common Defects**

44 Severely Damaged  
44 Broken, Chipped, Loose Seed Coat and Split  
44 Shrivelled and Wrinkled  
Sprouted  
Insect Damaged  
Poor Colour

### **46**

46  
46  
46  
46  
47  
47  
47  
48  
48  
48

### **50**

50  
50  
51  
51  
51  
52  
52  
52  
52

### **54**

54  
55  
56  
56  
56  
57



## **SORGHUM**

<b>Common Defects</b>	<b>59</b>
Severely Damaged	59
Insect Damaged	59
Sprouted	59
Mould	60
Field Fungi	60
Honeydew	61
Sorghum Ergot	61

## **WHEAT**

<b>Durum</b>	<b>63</b>
Durum Identification	63
Vitreous Kernels	63
<b>Common Defects</b>	<b>64</b>
Pink Stained	64
Stained	64
Insect Damaged	64
Distorted	65
Sprouted	65
Dry Green or Sappy	65
Field Fungi	66
Severely Damaged	66
White Grain Disorder	67
Ball Smuts	68
Pickling Compounds / Artificial Colouring (Pickled Wheat-WA)	68



# INTRODUCTION

## Defective Grains

This guide is produced to assist samplers and assessors of grain in the determination of defective grains which are covered by the Grain Trade Australia (GTA) and the Australian Oilseeds Federation (AOF) Standards.

**All images in this guide (unless otherwise identified) are defective. These photographs depict the minimum standard for a grain to be assessed as defective. If a grain defect does not meet the physical attributes depicted in the photograph, it is to be assessed as sound. Unless otherwise stated, the grain defect may only appear on one side. That is, the photographs depict the minimum to be assessed as defective, and only needs to be present on one side (unless otherwise stated).**




Pictures shown at this size are an approximate size of the original grain only.

These pictures are enlarged to assist in illustration of the defect.

## Calibration


A calibration sheet is provided for those who are downloading and printing these guides. Careful calibration of these photographs is vital as monitors and printers may vary.

# DOCUMENT CALIBRATION




CMYK Value:  
C=0 M=100 Y=0 K=0

Minolta Value:  
L= 48.59 a= +51.21 b= +31.27



CMYK Value:  
C=100 M=0 Y=0 K=0

Minolta Value:  
L= 55.41 a= -17.28 b= -43.99



CMYK Value:  
C=0 M=0 Y=100 K=0

Minolta Value:  
L= 87.53 a= -10.50 b= +80.56

NOTE: The hardware (monitor, graphics card, etc.) used to display the images in Inspector Standardisation content influences the appearance of the images. As a result the images may have a slightly different appearance when viewed on different makes/models of computer and display. These images were created using a Dino-Lite Edge 5MP AM7915, calibrated LCD display with 1680x1050, 32 Bit, 60 Hz resolution and the following calibration settings:

Brightness: 0

Contrast: 50

Gamma: 1.0

Hue: 0

Saturation: 0


The VRSG should be viewed using a computer with digital video (DVI) output and an EIZO CG19, EIZO S1921, EIZO S1932, EIZO S1961, or EIZO CE210W display.

Paper Type for Printing: Brand: Office Elements      GSM: 80gsm      Colour: White


Laminate material:      Brand: OfficeMax 125 Micron laminating pouches      Re-Order Code: 1950630

Disclaimer: The mention of firm names or trade products does not imply that they are endorsed or recommended.


# PRINTER CALIBRATION




CMYK Value:  
C=0 M=100 Y=0 K=0



CMYK Value:  
C=100 M=0 Y=0 K=0



CMYK Value:  
C=0 M=0 Y=100 K=0



CMYK Value:  
C=75 M=68 Y=0 K=100


NOTE: This page is specifically for use by a commercial printer.

Paper Type for Printing: Brand: Office Elements      GSM: 80gsm      Colour: White

Laminate material:      Brand: OfficeMax 125 Micron laminating pouches      Re-Order Code: 1950630

Disclaimer: The mention of firm names or trade products does not imply that they are endorsed or recommended.


# BARLEY



Barley Grain



Sound




# BARLEY: VARIETAL IDENTIFICATION

**Definition:** The main characteristic used in identifying barley varieties is the length of the hairs on the Rachilla. The Rachilla is white in colour and found running along the grain furrow from the germ end. There are two main types of Rachilla hair length, long hairs and short woolly hairs.



Long



Long

**Common Varieties include:**

- |             |             |
|-------------|-------------|
| Banks       | Maximus CL  |
| Bass        | Oxford      |
| Baudin      | Rosalind    |
| Beast (AGT) | Scope (Cl.) |
| Bottler     | Shepherd    |
| Commodus CL |             |
| Compass     |             |
| Fairview    |             |
| Fathom      |             |
| Grout       |             |
| Kiwi        |             |
| Leabrook    |             |

**Awn End of Grain**



**Dorsal (Back)**




**Ventral (Front)**

**Germ End of Grain**



Short



Short

**Common Varieties include:**

- |                 |                 |
|-----------------|-----------------|
| Buff            | Spartacus (Cl.) |
| Commander       | Urambie         |
| Fleet Australia | Westminster     |
| Flinders        |                 |
| Gardner         |                 |
| GrangeR         |                 |
| Hindmarsh       |                 |
| La Trobe        |                 |
| Laperouse       |                 |
| LG Alestar      |                 |
| (RGT) Planet    |                 |
| Schooner        |                 |



# BARLEY: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Sprouted

**Definition:** Sprouted grains are those with any visible evidence of the shoot or root system beginning to emerge from the germ.



Side

## Shot

**Definition:** Grains exhibiting the following outward signs of having commenced germination are classified as Shot:

- Opening of the grain at the germ end and/or
- The husk has 'tramlines' on both sides where the husk has begun to lift on each side on the back of the grain at the germ end.



"tramlines"

## Dark Tipped (Germ End Stained-WA)

**Definition:** Grains exhibit a distinct brown to black discoloration. This mainly occurs at the germ end of the grain, however in severe cases it may progress to other parts of the grain. Discolouration originating at the awn end is not Dark Tipped, refer to Severely Damaged. Dark tipping equal to or greater than 1mm is classified as defective grain.

Note: image represents the minimal level of intensity of colour and 1mm length.



Sound

Defective

# BARLEY: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Skinnings (Skinned-WA)

**Definition:** Damage to the protective husk of the barley. Usually caused by mechanical damage to the grain during harvesting.

Each grain exhibiting one or more of the following characteristics is assessed as a skinned grain -

- Skinning - 1/3 or more of the total surface area of the husk is missing.
- Germ Exposed - The husk is removed from the germ end of the grain or been damaged other than Shot or Sprouted or the germ itself has been removed.
- Kernels may or may not be dark under the husk.



Dorsal (Back)



Ventral (Front)



Germ Exposed



Dark Under Husk


## Cleaved (front, back and side)

**Definition:** Refers to any damage to the grain exposing the white endosperm. Any visible cleaving is considered defective. It includes but is not limited to -

- A split along the crease or a split down the back, front or side of the grain exposing the white endosperm.
- Grains that are swollen but smaller in size than normal, are sometimes fused in groups of 2 or 3 and are split to reveal the inner endosperm (sometimes identified as hormonal damage).
- Kernels may or may not have a pink discoloration.



Ventral (Front)



Ventral (Front)



Dorsal (Back)



Side



Hormonal Damage

# BARLEY: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024



## Distorted

**Definition:** Grain which is collapsed on the dorsal side. Grains may also appear orange in colour.



## Dry Green or Sappy

**Definition Dry Green:** Grain surface is distinctly green. Grains are usually dry and hard.


**Definition Sappy:** Grains are generally soft when pressed. They may or may not be green. Any level of sappiness is classified as defective.



Dry Green

## Broken

**Definition:** Grain that has 1/3 or more missing from the Kernel.



Broken

# BARLEY: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Heavily Discoloured-WA

**Definition:** Staining is dark in appearance. Usually affects more than the germ end.



## Pink Fungal Staining

**Definition:** Orange, pink to red discoloration found anywhere on the surface. Grains appear healthy otherwise.



## Field Fungi (Spotted/Field Fungi Affected-WA)

**Definition:** Refers to individual kernels -

- Seed coat has the appearance of black spotting occurring anywhere on the grain. Coverage greater than approximately 10% of the grain surface is considered defective, otherwise classified as sound.
- Grey or brown surface discoloration on the kernel and/or husk.



Black Spotting

Black Spotting

Grey Discolouration

Discolouration on Kernel



# BARLEY: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Severely Damaged inc. Fusarium (except WA)

**Definition:** Damage to the grain causing it to become severely discoloured. A grain exhibits one or more of the following characteristics -

### Burnt / Heat Damaged:

Heat Damaged or Burnt refers to those kernels that have become discoloured. Affected grains appear dark brown, or in severe cases, blackened. May also appear discoloured under the husk on the kernel.


### Mould:

Affected grains appear discoloured and visibly affected by mould.

### Diseased / Other Serious Visual Defects:

Refers to those kernels that have become significantly discoloured and/or have a serious visual defect that is not otherwise listed in these Standards. It includes grains affected by Fusarium (except WA).

Does not include Field Fungi affected grains, refer to Field Fungi.



# BARLEY: CONTAMINANTS/DEFECTS

Issued: 1<sup>st</sup> August 2024

## Coloured Aleurone Layer (Blue or Black)

**Definition:** Grains which have a coloured aleurone layer in the kernel. The colour is generally blue or black.

It includes any blue or black colour to any degree that is obvious under the bran layer.

Also includes black hulled varieties.



Blue Aleurone Layer



Black Hulled

## Insect Damaged

**Definition:** Any visible insect damage penetrating through to the white endosperm.



## Pickling Compounds or Artificial Colour (Pickled Barley-WA)

**Definition:** An unnatural surface colour and/or colour that rubs off. Any grains that are artificially coloured, regardless of intensity, are defective.


**Note:** These photographs are to illustrate artificial colours and appearance only. A **nil tolerance** applies to any pickling compounds/contaminants, regardless of intensity or coverage or colour.



# CANOLA



Sound



# CANOLA: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Broken or Split

**Definition:** All hulls, kernels or parts thereof, not otherwise damaged shall be classified as split or broken seed (except fines classified as Impurities). Any level of damage is classified as defective. This includes Insect Damaged.



## Heat Damaged

**Definition:** Seeds and pieces of seed that are materially discoloured and damaged by heat. Seeds may have a heated odour or a brown powdery appearance when crushed.



Sound Crushed



Defective Crushed


## Sprouted

**Definition:** The seed coat has split and the primary root has emerged. This includes early and any further advanced stage of growth of the primary root. Includes grains where the primary root has been knocked off during the harvesting or handling process. Seed has visibly swelled and enlarged.



## Weather Damaged

**Definition:** Weather damaged seeds are classified under Damaged Seeds. Weather damaged seeds are those that have been subjected to rain during the maturation phase to the extent they have become Weather Damaged. When seeds are crushed, they may have a grey washed out appearance and a chalky texture.



Sound Crushed



Defective Crushed

# CANOLA: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Mould

**Definition:** Seeds may appear discoloured, rotten, swollen and soft, feel spongy under pressure, show the presence of fungal spores or visibly affected by mould on the seed coat. Includes Field Fungi seeds.



## Frost Damaged

**Definition:** Frost damaged seeds are included in Impurities. Frost damaged seeds are included in admixture (WA).



## Green Seeds

**Definition:** Green seeds are those that are distinctly green when crushed. Seeds that are yellow-green are not considered green.



**Sound Crushed**  
Yellow or Yellow-Green when crushed




**Defective Crushed**  
Green Seed – distinctly green when crushed

# CHICKPEAS, DESI



Sound



# CHICKPEAS, DESI: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Severely Damaged

**Definition:** Damage to the grain causing it to become severely discoloured. A grain exhibits one or more of the following characteristics -

### Burnt / Heat Damaged:

Heat Damaged or Burnt refers to those grains that have become severely discoloured. Affected grains appear dark brown, or in severe cases, blackened.

### Mould:

Affected grains appear discoloured and visibly affected by mould.

### Diseased / Other Serious Visual Defects:

Refers to those grains that have become significantly discoloured and/or have a serious visual defect that is not otherwise listed in these Standards.



**Sound -  
Refer Stained and  
Weather Damaged.**



**Heat Damaged**



**Mould**



**Mould**

# CHICKPEAS, DESI: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Broken, Chipped, Loose Seed Coat and Split

**Definition:** Breakage, cracking, peeling or splitting of the seed coat or chipping and splitting of the kernel in various forms as follows -

### Seed Coat:

- Split Seed Coat - A split in the seed coat running more than half the entire length or across the entire width on one or both sides.
- Skin Damaged - A hole in the seed coat where more than 20% of the seed coat on any one side is missing.
- Loose Seed Coat (Peeling) - Where the seed coat is visibly falling off the kernel to any extent and not adhering tightly to the kernel.
- Missing Seed Coat - Where the entire seed coat is missing but the kernel remains intact.



### Kernel:

- Chipped (Scratched) – A part of the kernel is damaged or removed.
- Broken - A split kernel with the seed coat still attached.
- Split – A split kernel with no seed coat attached.



# CHICKPEAS, DESI: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Insect Damaged


**Definition:** Any visible insect damage to the grain is to be classified as defective. Includes insect eggs of all Stored Grain Insect species on the Seed Coat surface, whether those eggs may be live or dead and in any number.



Stored Grain Insect Egg

## Sprouted

**Definition:** The seed coat has split and the primary root has emerged. This includes early and any further advanced stage of growth of the primary root. Includes grains where the primary root has been knocked off.



## Hail Damaged

**Definition:** Any damage to the seed coat or kernel is classified as defective. Damage to the seed coat can appear as bruising (darkening) or in more severe cases splitting of the seed coat. This may cause discoloration and damage to the kernel. Damage to the kernel can vary from bruising (darkening) to physical damage such as crushing of the entire kernel.



# CHICKPEAS, DESI: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Shrivelled and Wrinkled

**Definition:** Visible damage to the seed coat or size and shape of grain whereby the grains are severely distorted and/or shrunken. A distinct ridge (often described as mountains and valleys) on the seed coat must be present to be classified as Shrivelled & Wrinkled. Ridges may be described as coarse waves rather than soft waves. Seed coats may be wrinkled or dimpled and distinctly indented into the kernel. Seed coats may show a level of discolouration depending on the extent of damage. Grains are often smaller than the majority in the sample.



## Green

**Definition:** Green is included in Poor Colour.


### Seed Coat:

Seed coat appears green. More than a slight greenish tinge must be present on the seed coat to be classified as defective.

Where any greenish tinge is present on the seed coat, it is recommended the kernel also be inspected.

### Kernel:

Any level of green is classified as defective.



Green Kernel

## Pickling Compounds or Artificial Colour

**Definition:** An unnatural surface colour and/or colour that rubs off. Any grains that are artificially coloured, regardless of intensity, are defective.

**Note:** These photographs are to illustrate artificial colours and appearance only. A **nil tolerance** applies to any pickling compounds/contaminants, regardless of intensity or coverage or colour.



# CHICKPEAS, DESI: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Poor Colour


**Definition:** Green is included in Poor Colour. Stained and Weather Damaged is included in Poor Colour. Fungal Affected is included in Poor Colour.

### Seed Coat:

Seed coats vary from dark brown to black, but may be depicted by other colours. Seed coats may be similar in appearance to various other defects such as Severely Damaged.

Where any poor colour is present on the seed coat, it is recommended the kernel also be inspected.

Poor Colour kernel can only be assessed if the seed coat is removed.




### Kernel:

Any level of discolouration on the kernel is classified as defective.



**Not Poor Colour Kernel.  
Refer Poor Colour Seed Coat.**



**Defective Kernel, therefore this is classified as Poor Colour.**



# CHICKPEAS, DESI: COMMON DEFECTS


Issued: 1<sup>st</sup> August 2024

## Stained and Weather Damaged

**Definition:** Stained and Weather Damaged is included in Poor Colour.

A general term used to describe visible damage to the seed coat that may or may not otherwise be defined or be distinguishable from other defects in the Standards. Seed coats may be discoloured or altered in size or shape. Weather Damage may also lead to a Loose Seed Coat or Shrivelled and Wrinkled.

Any degree of Tiger Striping or Speckled is not considered Poor Colour, but is considered sound. Tiger Striping or Speckling on the kernel is to be classified as Poor Colour Kernel.



## Fungal Affected (e.g. Ascochyta)

**Definition:** Fungal Affected is included in Poor Colour.

Lesions are generally visible to the naked eye and appear intense dark brown to black. The lesion may be similar in colour to Severely Damaged or Stained and Weather Damaged.


Any lesion of any size is permitted provided it is not also present on the kernel. If the lesion is greater than approximately 20%, but does not penetrate to the kernel the grain is classified as Stained and Weather Damaged.



# CHICKPEAS, KABULI



Sound



# CHICKPEAS, KABULI: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Poor Colour


**Definition:** Seed coats vary from dark brown to black, but may be depicted by other colours. Seed coats may be similar in appearance to various other defects such as Severely Damaged.

Where any poor colour is present on the seed coat, it is recommended the kernel also be inspected.



## Insect Damaged

**Definition:** Any visible insect damage to the grain is to be classified as defective. Includes insect eggs of all Stored Grain Insect species on the Seed Coat surface, whether those eggs may be live or dead and in any number.



# CHICKPEAS, KABULI: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Broken, Chipped, Loose Seed Coat and Split

**Definition:** Breakage, cracking, peeling or splitting of the seed coat or chipping and splitting of the kernel in various forms as follows -


### Seed Coat:

- Split Seed Coat - A split in the seed coat running more than half the entire length or across the entire width on one or both sides.
- Skin Damaged - A hole in the seed coat where more than 20% of the seed coat on any one side is missing.
- Loose Seed Coat (Peeling) - Where the seed coat is visibly falling off the kernel to any extent and not adhering tightly to the kernel.
- Missing Seed Coat - Where the entire seed coat is missing but the kernel remains intact.



### Kernel:

- Chipped (Scratched) – A part of the kernel is damaged or removed.
- Broken - A split kernel with the seed coat still attached.
- Split – A split kernel with no seed coat attached



# CHICKPEAS, KABULI: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Severely Damaged

**Definition:** Damage to the grain causing it to become severely discoloured. A grain exhibits one or more of the following characteristics:

### Burnt / Heat Damaged:

Heat Damaged or Burnt refers to those grains that have become severely discoloured. Affected grains appear dark brown, or in severe cases, blackened.

### Mould:

Affected grains appear discoloured and visibly affected by mould.

### Diseased / Other Serious Visual Defects:

Refers to those grains that have become significantly discoloured and/or have a serious visual defect that is not otherwise listed in these Standards.




**Mould**

**Mould**

## Sprouted

**Definition:** The seed coat has split and the primary root has emerged. This includes early and any further advanced stage of growth of the primary root. Includes grains where the primary root has been knocked off.



## Shriveled and Wrinkled

**Definition:** Visible damage to the seed coat or size and shape of grain whereby the grains are severely distorted and/or shrunken. A distinct ridge (often described as mountains and valleys) on the seed coat must be present to be classified as Shriveled & Wrinkled. Ridges may be described as coarse waves rather than soft waves. Seed coats may be wrinkled or dimpled and distinctly indented into the kernel. Seed coats may show a level of discoloration depending on the extent of damage. Grains are often smaller than the majority in the sample.



**Sound**

**Defective**

**Defective**

# FABA BEANS

*\*\*Where applicable, Broad Beans should be assessed using Faba Bean visual images.*



Sound



# FABA BEANS: COMMON DEFECTS

## Severely Damaged

**Definition:** Damage to the grain causing it to become severely discoloured. A grain exhibits one or more of the following characteristics -

### Burnt / Heat Damaged:

Heat Damaged or Burnt refers to those grains that have become severely discoloured. Affected grains appear dark brown, or in severe cases, blackened.

### Mould:

Affected grains appear discoloured and visibly affected by mould.

### Diseased / Other Serious Visual Defects:

Refers to those grains that have become significantly discoloured and/or have a serious visual defect that is not otherwise listed in these Standards.



**Pod Fluff  
Sound**



**Burnt**



**Heat Damaged**



**Heat Damaged**



**Other Serious  
Visual Defect**



**Mould**




**Mould**

# FABA BEANS: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Shrivelled and Wrinkled


**Definition:** Visible damage to the seed coat or size and shape of grain whereby the grains are severely distorted and/or shrunken. A distinct ridge (often described as mountains and valleys) on the seed coat must be present to be classified as Shrivelled & Wrinkled. Ridges may be described as coarse waves rather than soft waves. Seed coats may be wrinkled or dimpled and distinctly indented into the kernel. Seed coats may show a level of discolouration depending on the extent of damage. Grains are often smaller than the majority in the sample.



## Sprouted

**Definition:** Seed coat has split and the primary root has emerged. This includes early and any further advanced stage of growth of the primary root. Includes grains where the primary root has been knocked off.

Where the seed coat has split but the primary root has not emerged, the grain is to be classified under Broken, Chipped, Loose Seed Coat and Split.



## Insect Damaged

**Definition:** Any visible insect damage to the grain is to be classified as defective. Includes insect eggs of all Stored Grain Insect species on the Seed Coat surface, whether those eggs may be live or dead and in any number.



# FABA BEANS: COMMON DEFECTS

## Broken, Chipped, Loose Seed Coat and Split

**Definition:** Breakage, cracking, peeling or splitting of the seed coat or chipping and splitting of the kernel in various forms as follows -


### Seed Coat:

- Split Seed Coat - A split in the seed coat running more than half the entire length or across the entire width on one or both sides. Split may or may not be tightly adhering to the kernel.
- Skin Damaged - A hole in the seed coat where more than 20% of the seed coat on any one side is missing.
- Loose Seed Coat (Peeling) - Where the seed coat is visibly falling off the kernel to any extent and not adhering tightly to the kernel.
- Missing Seed Coat - Where the entire seed coat is missing but the kernel remains intact.



### Kernel:

- Chipped (Scratched) – A part of the kernel is damaged or removed.
- Broken - A split kernel with the seed coat still attached.
- Split – A split kernel with no seed coat attached.



# FABA BEANS: COMMON DEFECTS

## Poor Colour

**Definition:** Green is included in Poor Colour. Fungal Affected is included in Poor Colour. Frost Damaged, Stained is included in Poor Colour. Pea Seed Borne Mosaic Virus is included in Poor Colour.

Seed coats vary from grey, dark brown to black but may be depicted by other colours. Seed coats may be similar in appearance to various other defects such as Severely Damaged.

The photos below depict the minimum requirement of any colour to be classified as defective.



Defective

## Green


**Definition:** Green is included in Poor Colour.

### Seed Coat:

Seed coat appears intense green. It is recommended the kernel be inspected if immature grains are present.

### Kernel:

Any level of green is classified as defective.



# FABA BEANS: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Fungal Affected (e.g. Ascochyta)

**Definition:** Fungal Affected is included in Poor Colour.

Lesions are generally visible to the naked eye and appear intense dark brown to black. The lesion may be similar in colour to Severely Damaged or Stained and Weather Damaged. A lesion may appear on one or both sides of the seed coat or kernel.

A lesion greater than 20% coverage on any one side of the seed coat is considered defective.

Any lesion of any size on the kernel is defective.



## Stained and Weather Damaged

**Definition:** Stained and Weather Damaged is included in Poor Colour.

A general term used to describe visible damage to the seed coat that may or may not otherwise be defined or be distinguishable from other defects in these Standards. Weather Damage may also lead to a Loose Seed Coat or Shrivelled and Wrinkled.

### Seed Coat:

Visible damage to the seed coat resulting in staining on the seed coat only. Seed coats may be discoloured or altered in size or shape.

### Kernel:

Any damage to the kernel is classified as defective.



**Defective**  
(Seed Coat Affected)

**Defective (Kernel Affected)**  
Photos do not depict minimum required

# FABA BEANS: COMMON DEFECTS

## Pea Seed Borne Mosaic Virus

**Definition:** Pea Seed Borne Mosaic Virus is included in Poor Colour. Staining on the seed coat caused by the Pea Seed Borne Mosaic Virus.



# LENTILS, RED




**Whole Seed  
Sound**



**Kernel  
Sound**

Note: Green Lentils are considered a contaminant in Red Lentils.



# LENTILS, RED: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Shrivelled and Wrinkled

**Definition:** Visible damage to the seed coat or size and shape of grain whereby the grains are severely distorted and/or shrunken. A distinct ridge (often described as mountains and valleys) on the seed coat must be present to be classified as Shrivelled & Wrinkled. Ridges may be described as coarse waves rather than soft waves. Seed coats may be wrinkled or dimpled and distinctly indented into the kernel. Seed coats may show a level of discolouration depending on the extent of damage. Grains are often smaller than the majority in the sample.



# LENTILS, RED: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Broken, Chipped, Loose Seed Coat and Split

**Definition:** Breakage, cracking, peeling or splitting of the seed coat or chipping and splitting of the kernel in various forms as follows -

### Seed Coat:

- Split Seed Coat - A split in the seed coat running more than half the entire width or across the entire width on one or both sides.
- Skin Damaged - A hole in the seed coat where more than 20% of the seed coat on any one side is missing.
- Loose Seed Coat (Peeling) - Where the seed coat is visibly falling off the kernel to any extent and not adhering tightly to the kernel.
- Missing Seed Coat - Where the entire seed coat is missing but the kernel remains intact.



### Kernel:

- Chipped (Scratched) – A part of the kernel is damaged or removed.
- Broken - A split kernel with the seed coat still attached.
- Split – A split kernel with no seed coat attached




# LENTILS, RED: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024


## Insect Damaged

**Definition:** Any visible insect damage to the grain is to be classified as defective. Includes insect eggs of all Stored Grain Insect species on the Seed Coat surface, whether those eggs may be live or dead and in any number.



## Sprouted

**Definition:** Seed coat has split and the primary root has emerged. This includes early and any further advanced stage of growth of the primary root. Includes grains where the primary root has been knocked off.



## Severely Damaged

**Definition:** Damage to the grain causing it to become severely discoloured. A grain exhibits one or more of the following characteristics -

### Burnt / Heat Damaged:

Heat Damaged or Burnt refers to those grains that have become severely discoloured. Affected grains appear dark brown, or in severe cases, blackened.

### Mould:

Affected grains appear discoloured and visibly affected by mould.

### Diseased / Other Serious Visual Defects:

Refers to those grains that have become significantly discoloured and/or have a serious visual defect that is not otherwise listed in these Standards.



# LENTILS, RED: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Poor Colour (Seed Coat)

**Definition:** Fungal Affected is included in Poor Colour. Stained and Weather Damaged is included in Poor Colour.

Seed coats vary from dark brown to black, but may be depicted by other colours. Seed coats may be similar in appearance to various other defects such as Severely Damaged. Does not include Contrasting Colour.


Where any poor colour is present on the seed coat, it is recommended the kernel also be inspected. Poor Colour kernel can only be assessed if the seed coat is removed.



Defective


## Orange Tip

**Definition:** Seed coats have varying degrees of orange colour, generally in the germ area. Included in defective not Poor Colour.



## Poor Colour Kernel - Dehulled Lentil

**Definition:** Poor Colour Kernel refers to excessive discolouration of the kernel often depicted as a green colour. Includes green, brown, black, yellow, bleached and chalky white kernels or any other discolouration.



Sound

Poor Colour Kernel

Poor Colour Kernel

Poor Colour Kernel

## Blonde Kernel

**Definition:** Kernels are not uniformly orange in colour. Kernels appear yellow. Seed coat must be removed to determine the presence on the kernel. Included in defective not Poor Colour.



# LENTILS, RED: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Fungal Affected

**Definition:** Fungal Affected is included in Poor Colour.

Lesions are generally visible to the naked eye and appear intense dark brown to black. The lesion may be similar in colour to Severely Damaged or Stained and Weather Damaged.

A lesion greater than 20% coverage on any one side of the seed coat is considered defective.

Any lesion of any size on the kernel is defective.



Defective

## Stained and Weather Damaged


**Definition:** Stained and Weather Damaged is included in Poor Colour.

A general term used to describe visible damage to the seed coat or kernel that may or may not otherwise be defined or be distinguishable from other defects in these Standards. Seed coats and kernels may be discoloured or altered in size or shape. Weather damage may also lead to Loose Seed Coat or Shrivelled and Wrinkled.

Does not include Contrasting Colour.



Speckled -  
Sound



Defective

# LENTILS, RED: CONTRASTING COLOURS

Issued: 1<sup>st</sup> August 2024



## Variety Definition Chart - Supplied by Grains Australia

**Definition:** Grains with a colour not falling within the 'main and acceptable variety seed coat variation' are to be classified as Contrasting Colour.

Contrasting colour and marbling is genetic variation, which can occur at trace levels within specific varieties and are not defective grains.

Contrasting Colour	Main and acceptable variety seed coat variation – Sound					Contrasting Colour
<b>PBA Blitz</b> <sup>Ⓛ</sup>  Pale	 Typical Grey	 Grey-green	 Slightly Marbled	 Medium Marbled	 Strongly Marbled	
<b>PBA Herald XT</b> <sup>Ⓛ</sup>  Pale	 Typical Grey	 Grey-green	 Slightly Marbled	 Medium Marbled	 Strongly Marbled	 Black (totally marbled)
<b>PBA Hurricane XT</b> <sup>Ⓛ</sup>  Pale	 Typical Grey	 Grey-green				
<b>GIA Lightning</b> <sup>Ⓛ</sup> and <b>GIA Thunder</b> <sup>Ⓛ</sup>	 Typical Grey	 Grey-green	 Slightly Marbled	 Medium Marbled	 Strongly Marbled	 Black (totally marbled)

# LUPINS, ANGUSTIFOLIUS




**Lupin -  
Angustifolius  
Sound**



**Lupin -  
Albus  
Sound**

Note: Albus Lupins are considered a contaminant in Angustifolius Lupins.



# LUPINS, ANGUSTIFOLIUS: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Broken, Chipped, Loose Seed Coat and Split

**Definition:** Breakage, cracking, peeling or splitting of the seed coat or chipping and splitting of the kernel in various forms as follows -


### Seed Coat:

- Split Seed Coat - A split in the seed coat running more than half the entire length or across the entire width on one or both sides.
- Skin Damaged - A hole in the seed coat where more than 20% of the seed coat on any one side is missing.
- Loose Seed Coat (Peeling) - Where the seed coat is visibly falling off the kernel to any extent and not adhering tightly to the kernel.
- Missing Seed Coat (Fully De-Coated - WA) - Where the entire seed coat is missing but the kernel remains intact.



### Kernel:

- Chipped (Scratched) – A part of the kernel is damaged or removed.
- Broken - A split kernel with the seed coat still attached.
- Split – A split kernel with no seed coat attached



## Manganese Deficiency

**Definition:** Splitting of the seed coat to expose the kernel. It is not considered a defect provided no damage to the exposed kernel has occurred.




Sound

# LUPINS, ANGUSTIFOLIUS: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Shriveled and Wrinkled

**Definition:** Visible damage to the seed coat or size and shape of grain whereby the grains are severely distorted and/or shrunken. A distinct ridge (often described as mountains and valleys) on the seed coat must be present to be classified as Shriveled & Wrinkled. Ridges may be described as coarse waves rather than soft waves. Seed coats may be wrinkled or dimpled and distinctly indented into the kernel. Seed coats may show a level of discoloration depending on the extent of damage. Grains are often smaller than the majority in the sample.



## Insect Damaged

**Definition:** Any visible insect damage to the grain is to be classified as defective. Includes insect eggs of all Stored Grain Insect species on the Seed Coat surface, whether those eggs may be live or dead and in any number.




# LUPINS, ANGUSTIFOLIUS: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Poor Colour (Discoloured-WA)

**Definition:** Seed coats vary from yellow to tan, dark brown to black, but may be depicted by other colours. Seed coats may be similar in appearance to various other defects such as Severely Damaged.



## Phomopsis

**Definition:** Grains appear sound with a fungal growth readily visible on the seed coat. If kernels are not sound, refer to Severely Damaged.



## Bitter Dark (Bitter Variety-WA)

**Definition:** These varieties are identifiable mainly by their colour which is much darker than acceptable lupins.



# LUPINS, ANGUSTIFOLIUS: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Severely Damaged

**Definition:** Damage to the grain causing it to become severely discoloured. A grain exhibits one or more of the following characteristics -

### Burnt / Heat Damaged:

Heat Damaged or Burnt refers to those grains that have become severely discoloured. Affected grains appear dark brown, or in severe cases, blackened.

### Mould:

Affected grains appear discoloured and visibly affected by mould. Does not include Phomopsis.

### Diseased / Other Serious Visual Defects:

Refers to those grains that have become significantly discoloured and/or have a serious visual defect that is not otherwise listed in these Standards.



Heat Damaged /  
Burnt

Mould

## Sprouted

**Definition:** Seed coat has split and the primary root has emerged. This includes early and any further advanced stage of growth of the primary root. Includes grains where the primary root has been knocked off.



## Pickling Compounds or Artificial Colour

**Definition:** An unnatural surface colour and/or colour that rubs off. Any grains that are artificially coloured, regardless of intensity, are defective.

**Note:** This photograph is to illustrate artificial colour and appearance only. A **nil tolerance** applies to any pickling compounds, regardless of intensity or coverage or colour.



# MAIZE, FEED



Sound



# MAIZE, FEED: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Heat Damaged / Bin Burnt

**Definition:** Grains appear reddish brown, or in severe cases, blackened. Heat Damaged / Bin Burnt is included in the definition of Damaged.



## Insect Damaged

**Definition:** Any visible insect damage to the grain is to be classified as defective. Insect Damaged is included in the definition of Damaged.



## Broken

**Definition:** Grains that have 1/4 or more missing from the Kernel.



## Sprouted

**Definition:** Included in Damaged.

Grains in which the covering of the germ is split and the shoot has broken through the seed coat. Grains that have had the germ knocked off or scalloped.



# MAIZE, FEED: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Storage Mould

**Definition:** Grains appear discoloured and visibly affected by mould.

**Note:** If any musty odour is detected a nil tolerance applies.



## Dead

**Definition:** Grains that are at least 50% opaque.



## Fungal Affected

### Definitions:

#### Silk Cut:

Easily identified where the pericarp is split and the starch appears to be popping out of the kernel.

Silk Cut is included in Damaged.

#### Starburst:

Best identified as spider web like streaks radiating down the kernel from the point of silk attachment.

These streaks are corroded channels within the pericarp caused by fungal growth. Air in the channels breaks the transparency of the pericarp so the yellow aleurone beneath cannot be seen.

Star Burst is included in the definition of Dead, Mouldy, Storage Mould.



Silk Cut

Star Burst

# MAIZE, FEED: COMMON DEFECTS/CONTAMINANTS

## Pickling Compounds or Artificial Colouring

**Definition:** An unnatural surface colour and/or colour that rubs off. Any grains that are artificially coloured, regardless of intensity, are defective.


**Note:** This photograph is to illustrate artificial colour and appearance only. A **nil tolerance** applies to any pickling compounds, regardless of intensity or coverage or colour.



## Field Fungi

**Definition:** Included in Damaged.

Seed coat is greater than approximately 50% discoloured. The visible discolouration of affected grains can vary from dark grey, brown to black in colour.




## Kernel Red Streak

**Definition:** This is not a defect and kernels are considered sound when identified as Kernel Red Streak.



Sound

# OATS



Awn End of Grain



Groatsound

Oatsound

Germ End of Grain



# OATS: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024



## Severely Damaged

**Definition:** Damage to the grain causing it to become severely discoloured. A grain exhibits one or more of the following characteristics -

### Burnt / Heat Damaged:

Heat Damaged or Burnt refers to those kernels that have become discoloured. Affected grains appear dark brown, or in severe cases, blackened. May also appear discoloured under the husk on the groat.

### Mould:

Affected grains appear discoloured and visibly affected by mould. Note that light Septoria discolouration similar to Mould is not included in the definition of Severely Damaged - refer to Stained Grains.

### Diseased / Other Serious Visual Defects:

Refers to those kernels that have become significantly discoloured and / or have a serious visual defect that is not otherwise listed in these Standards. Does not include Field Fungi affected grains, refer to Field Fungi.



**Sound -  
Variety differs**

**Heat damaged**

## Field Fungi (Spotted Mould Affected-WA)

**Definition:** Individual kernels where the seed coat has grey to black spotting occurring anywhere on the grain.

Coverage greater than approximately 10% of the grain surface is considered defective, otherwise classified as sound.

Grains that are soft (that are not classified as Sappy) and/or emit a mouldy odour are to be classified as Objectionable Material.



**Dorsal**

**Dorsal**

**Ventral**

**Ventral**

# OATS: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Stained Grains (Heavily Discoloured-WA) inc. Septoria

**Definition:** Grains where greater than approximately 50% of the grain surface is discoloured. Various colours may be exhibited such as brown to black. Septoria is to be included in this definition, except in WA.


Grains that are affected by Field Fungi or Mould are not included in the definition of Stained Grains.

Where staining has occurred, it is recommended the husk be removed and the Groat examined.



## Septoria (WA)

**Definition:** A fungal infection that causes light to dark discoloration on the husk and/or groat.



Side

## Stained Groats (Except WA)

**Definition:** This defect is checked where Stained Grains are present in the sample. Where this staining has occurred, it is recommended the husk be removed and the Groat examined.

Any discolouration from the normal colour of the Groat is defective.



Sound - Dorsal

Sound - Ventral

Stained - Side

Stained - Dorsal

Stained - Dorsal

Stained - Ventral

# OATS: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Insect Damaged

**Definition:** Any visible insect damage to the grain is to be classified as defective.



## Sprouted

**Definition:** The grain has begun the germination process. A kernel that is sprouted is one where the shoot is visibly seen growing out from the germ.



## Damaged Grains


**Definition:** Grain with a quarter or more of the grain missing. This includes any mechanical damage to the germ.



## Dry Green or Sappy

**Definition Dry Green:** Grain surface is distinctly green. Grains are usually dry and hard.


**Definition Sappy:** Grains are generally soft when pressed. They may or may not be green. Any level of sappiness is classified as defective.



Dry Green

## Shot

**Definition:** The covering of the germ is split, but without further development of the shoot.



# PEAS, FIELD



Typical White type Pea



Typical Blue type Pea



Sound



Acceptable colour range for Parafield type Peas



Typical shape for Parafield type Peas



Typical shape for Kaspas type Peas



Acceptable colour range for Kaspas type Peas



# PEAS, FIELD: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Severely Damaged

**Definition:** Damage to the grain causing it to become severely discoloured. A grain exhibits one or more of the following characteristics -

### Burnt / Heat Damaged:

Heat Damaged or Burnt refers to those grains that have become severely discoloured. Affected grains appear dark brown, or in severe cases, blackened.

### Mould:

Affected grains appear discoloured and visibly affected by mould.

### Diseased / Other Serious Visual Defects:

Refers to those grains that have become significantly discoloured and/or have a serious visual defect that is not otherwise listed in these Standards.



**Burnt / Heat  
Damaged**



**Mould**

# PEAS, FIELD: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Broken, Chipped, Loose Seed Coat and Split

**Definition:** Breakage, cracking, peeling or splitting of the seed coat or chipping and splitting of the kernel in various forms as follows -

### Seed Coat:

- Split Seed Coat - A split in the seed coat running more than half the entire width or across the entire width on one or both sides.
- Skin Damaged - A hole in the seed coat where more than 20% of the seed coat on any one side is missing.
- Loose Seed Coat (Peeling) - Where the seed coat is visibly falling off the kernel to any extent and not adhering tightly to the kernel.
- Missing Seed Coat - Where the entire seed coat is missing but the kernel remains intact.



### Kernel:

- Chipped (Scratched) - A part of the kernel is damaged or removed.
- Broken - A split kernel with the seed coat still attached.
- Split - A split kernel with no seed coat attached.



# PEAS, FIELD: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Shrivelled and Wrinkled

**Definition:** Visible damage to the seed coat or size and shape of grain whereby the grains are severely distorted and/or shrunken. A distinct ridge (often described as mountains and valleys) on the seed coat must be present to be classified as Shrivelled & Wrinkled. Ridges may be described as coarse waves rather than soft waves. Seed coats may be wrinkled or dimpled and distinctly indented into the kernel. Seed coats may show a level of discolouration depending on the extent of damage. Grains are often smaller than the majority in the sample.



## Sprouted

**Definition:** Seed coat has split and the primary root has emerged. This includes early and any further advanced stage of growth of the primary root. Includes grains where the primary root has been knocked off.



## Insect Damaged

**Definition:** Any visible insect damage to the grain is to be classified as defective. Includes insect eggs of all Stored Grain Insect species on the Seed Coat surface, whether those eggs may be live or dead and in any number.



# PEAS, FIELD: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024


## Poor Colour

### Definition:

### Seed Coat:

Seed coats vary from dark brown to black but may be depicted by other colours. Seed coats may be similar in appearance to various other defects such as Severely Damaged.

Where poor colour is present on the seed coat, it is recommended the kernel also be inspected.



### Kernel:

Any level of discolouration on the kernel is classified as defective. Where green kernels exist, the level of green colouring classified as defective is shown in the photos below.



# SORGHUM



Sound



# SORGHUM: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Severely Damaged


**Definition:** Heat Damaged / Burnt, diseased or other serious visual defects.

Kernels have become significantly discoloured. Grains appear dark brown or in severe cases, blackened.



## Insect Damaged

**Definition:** Any visible insect damage to the grain is to be classified as defective.



## Sprouted

**Definition:** The shoot is visibly extending from any part of the germ. Grains with pin holes are not included in this definition.


Scalloped grains are included in this definition.



Sound



Sprouted




Scalloped

# SORGHUM: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024


## Mould

**Definition:** Grains appear discoloured and visibly affected by mould.




## Field Fungi

**Definition:** Kernels affected by the growth of fungi on the seed coat. The fungal growth can vary in colour from white, to grey, to black. It does not refer to the more serious Moulds.



**Stained  
Sound**



**Field Fungi  
Defective**




# SORGHUM: COMMON DEFECTS/CONTAMINANTS

Issued: 1<sup>st</sup> August 2024

## Honeydew

**Definition:** Honeydew is acceptable if the grain is able to flow freely.

Honeydew is a sticky exudate produced by the sorghum plant in response to any predator attack, including Ergot. Honeydew oozes out of the flowers and drips onto leaves of the sorghum plant. It causes seeds to stick together and can make crops difficult to harvest and prevent harvested grain from running through equipment.




## Sorghum Ergot

**Definition:** Any visible ergot to the grain is to be classified as defective.

Sorghum Ergot, *Claviceps africana*, may result in the accumulation of a grey/white fungal mass in empty seed glumes. *Cerebella spp.* is not a true ergot but is a fungus that often grows on the *Claviceps africana*, producing a large black mass. *Cerebella spp.* is included in Sorghum Ergot.




**Claviceps africana**




**Cerebella spp.**

# WHEAT



Wheat Grain

Brush End of Grain




White Wheat Sound



Red Wheat Sound

Germ End of Grain




# WHEAT: DURUM

Issued: 1<sup>st</sup> August 2024

## Durum Identification

**Description:** Bread Wheat can be visually distinguished from Durum by the “fine hairs” on the brush end of the grain on some varieties. While these brush end hairs are usually only associated with bread wheat varieties, some durum varieties may also have brush end hairs of varying length.



## Vitreous Kernels

**Description:** Vitreous grains will appear uniformly bright and translucent. Non vitreous grains will be dull and opaque, or will contain dark, opaque sections that are clearly visible within the remaining translucent section of the grain.

Bleached grains may be difficult to ascertain if they are vitreous.

Any level of non-vitreous results in the grain being classified as non-vitreous.



Any level of non-vitreous results in non-vitreous classification.


# WHEAT: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Pink Stained

**Definition:** Grains with distinct pink discolouration.

Grains that are pink but also contain a white to light grey fungal like discolouration over more than approximately 50% of the seed coat surface are to be classified as "White Grain Disorder".



## Insect Damaged

**Definition:** Any visible insect damage to the grain is classified as defective.



## Stained

**Definition:** A distinct dark brown to black discolouration on the germ end that, in severe cases, may progress to other parts of the grain such as the crease. Grains are commonly referred to as "black point" or "black tip". Includes any staining beyond the minimum and up to 50% of the entire grain surface. For staining level greater than 50%, refer to Field Fungi.

Discolouration must be 50% or more of the germ

Includes grains that show streaking anywhere on the surface of the grain, and brush-end staining beyond the minimum. Also includes adherence of contaminants such as soil, dust, plant parts and other material.



Plant Material

Streaking

Brush Ventral

Brush Dorsal

Stained Crease

Black Tip

# WHEAT: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024



## Distorted

**Definition:** Grains generally have the appearance of full sized kernels with little or no structure on both dorsal sides of the grain, and are typically grey to blue in colour. Does not include pinched grain.



## Sprouted

**Definition:** The covering of the germ is split. It includes early and any further advanced stage of growth of the germ. Kernels exhibiting early stages of sprouting are those where the covering of the germ is split, but without further development of the shoot.

Grains with pin holes are not included in this definition.



Sound - Pin Hole

Scalloped  
Not defective

## Dry Green or Sappy

**Definition Dry Green:** Grain surface is distinctly green. Grains are usually dry and hard.

**Definition Sappy:** Grains are generally soft when pressed. They may or may not be green. Any level of sappiness is classified as defective.



Dry Green

# WHEAT: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## Field Fungi

**Definition:** Individual kernels where more than half (50%) of the seed coat is discoloured. Discolouration can vary from dark grey, brown to black in colour.

Grains that are approximately 50% or less discoloured are to be classified as Stained.



## Severely Damaged

**Definition:** Damage to the grain causing it to become severely discoloured. A grain exhibits one or more of the following characteristics -

### Burnt / Heat Damaged:

Heat Damaged or Burnt refers to those grains that have become severely discoloured. Affected grains appear reddish brown, dark brown or in severe cases, blackened.

### Mould:

Affected grains appear discoloured and visibly affected by mould.

### Diseased / Other Serious Visual Defects:

Refers to those grains that have become significantly discoloured and/or have a serious visual defect that is not otherwise listed in these Standards.



Heat Damaged

Heat Damaged  
(Blackened)

Mould

Other Serious  
Visual Defect

# WHEAT: COMMON DEFECTS

Issued: 1<sup>st</sup> August 2024

## White Grain Disorder

**Definition:** Includes Fusarium Head Blight. Grains appear white to light grey but may also contain a pink discoloration. Grains are only to be classified as White Grain Disorder if the discoloration is over more than approximately 50% of the seed coat surface. If the discoloration is approximately 50% or less of the seed coat surface, grains may be classified as Sound.

Grains may also appear flaky with a white discoloration and may display some level of shrivelling.



# WHEAT: COMMON DEFECTS/CONTAMINANTS

Issued: 1<sup>st</sup> August 2024

## Ball Smuts

**Definition:** Grains infected by the spores of the fungus *Tilletia caries*. Appearance of pale, plump, slightly oversized grains. Easily crushed between the fingers and contain a mass of black powder (spores) with a distinctive rotten egg smell. This may also be called Stinking Smut or Bunt.



## Pickling Compounds / Artificial Colouring (Pickled Wheat-WA)

**Definition:** An unnatural surface colour and/or colour that rubs off. Any grains that are artificially coloured, regardless of intensity, are defective.

**Note:** These photographs are to illustrate artificial colours and appearance only. A **nil tolerance** applies to any pickling compounds, regardless of intensity or coverage.





# GRAIN TRADE AUSTRALIA

Postal Address:

PO Box R1829

Royal Exchange NSW 1225

Email: [admin@graintrade.org.au](mailto:admin@graintrade.org.au)

Web: [www.graintrade.org.au](http://www.graintrade.org.au)