



GRAIN TRADE AUSTRALIA

Section 4 – AUSTRALIAN PULSE TRADING STANDARDS

2025/26 SEASON

TABLE OF CONTENTS

SECTION 1	INTRODUCTION	4
SECTION 2	DEFINITIONS	5
SECTION 3	GRAIN QUALITY STANDARDS	26
CSP-1.1	Adzuki Beans Minimum Receival Farmer Dressed.....	27
CSP-1.2	Adzuki Beans Minimum Export Standard Machine Dressed	28
CSP-2.1.1	Broad Bean Minimum Receival Farmer Dressed.....	29
CSP-2.1.2	Broad Beans Minimum Export Farmer Dressed	30
CSP-2.1.3	Broad Beans Minimum Export Machine Dressed	31
CSP-2.2	Broad Beans No.1 Split Minimum Export Standard.....	32
CSP-3.1	Caloona / Poona Cowpeas Minimum Export Standard Machine Dressed	33
CSP-4.1.1	Chickpea - Desi No.1 Minimum Receival Standard Farmer Dressed	34
CSP-4.1.2	Chickpea - Desi No.1 Minimum Export Standard Farmer Dressed.....	35
CSP-4.1.3	Chickpea - Desi No.1 Minimum Export Standard Machine Dressed.....	36
CSP-4.1.4	Chickpea - Desi No.2 Minimum Receival Standard Farmer Dressed	37
CSP-4.1.5	Chickpea - Desi No.2 Minimum Export Standard Farmer Dressed	38
CSP-4.1.6	Chickpea - Desi No.3 Minimum Receival Standard Farmer Dressed	39
CSP-4.1.7	Chickpea - Desi No.3 Minimum Export Standard Farmer Dressed	40
CSP-4.2	Chickpeas – Split Chana Dhal Minimum Export Standard.....	41
CSP-4.3.1	Chickpea Kabuli No.1 Large Minimum Receival Standard Farmer Dressed.....	42
CSP-4.3.2	Chickpea Kabuli No. 1 Large Minimum Export Standard Machine Dressed.....	43
CSP-4.3.3	Chickpea – Kabuli No.1 Small Minimum Receival Standard Farmer Dressed.....	44
CSP-4.3.4	Chickpea - Kabuli No.1 Small Minimum Export Standard Farmer Dressed	45
CSP-4.3.5	Chickpea – Kabuli No.1 Small Minimum Export Standard Machine Dressed	46
CSP-5.1.1	Faba Beans – Canning Grade Minimum Receival Standard Farmer Dressed.....	47
CSP-5.1.2	Faba Beans – Canning Grade Minimum Export Standard Machine Dressed.....	48
CSP-5.2.1	Faba Beans No.1 Minimum Receival Farmer Dressed.....	49
CSP-5.2.2	Faba Beans - No.1 Minimum Export Standard Farmer Dressed	50
CSP-5.2.3	Faba Beans No.1 Minimum Export Standard Machine Dressed	51

CSP-5.3.1 Faba Beans - No.2 Minimum Receival Standard Farmer Dressed 52

CSP-5.3.2 Faba Beans - No.2 Minimum Export Standard Farmer Dressed..... 53

CSP-5.4.1 Faba Beans - No.3 Minimum Receival Standard Farmer Dressed 54

CSP-5.5 Faba Beans – No.1 Split Minimum Export Standard Machine Dressed55

CSP-6.1 Fenugreek - Whole – No.1 Minimum Receival Standard Farmer Dressed..... 56

CSP-6.2 Fenugreek - Whole – No.1 Minimum Export Standard Machine Dressed.....57

CSP-7.1.1 Lentils – Whole Green No.1 Minimum Receival Standard Farmer Dressed 58

CSP-7.1.2 Lentils – Whole Green No.1 Minimum Export Standard Farmer Dressed..... 59

CSP-7.1.3 Lentils – Whole Green No.1 Minimum Export Standard Machine Dressed 60

CSP-7.2.1 Lentil - Whole Red No.1 Minimum Receival Standard Farmer Dressed61

CSP-7.2.2 Lentils – Whole Red No.1 Minimum Export Standard Farmer Dressed..... 62

CSP-7.2.3 Lentils – Whole Red No.1 Minimum Export Standard Machine Dressed..... 63

CSP-7.3.1 Lentils - Whole Red No.2 Minimum Receival Standard Farmer Dressed 64

CSP-7.3.2 Lentils Whole Red No.2 Minimum Export Standard Farmer Dressed..... 65

CSP-7.3.3 Lentil Whole Red No.2 Minimum Export Standard Machine Dressed 66

CSP-7.4.1 Lentils Split Red No.1 Minimum Export Standard67

CSP-7.4.2 Lentils Split Red No.2 Minimum Export Standard 68

CSP-7.4.3 Lentils Split Red No.3 Minimum Export Standard 69

CSP-8.1.1 Lupins Angustifolius Minimum Receival Standard Farmer Dressed 70

CSP-8.1.3 Lupins Angustifolius Minimum Export Standard Farmer Dressed 71

CSP-8.1.4 Lupins Angustifolius Minimum Export Standard Machine Dressed72

CSP-8.2.1 Lupins – Albus No.1 Minimum Receival Standard Farmer Dressed73

CSP-8.2.2 Lupins – Albus No.1 Minimum Export Standard Farmer Dressed.....74

CSP-8.2.3 Lupins – Albus No.1 Minimum Export Standard Machine Dressed.....75

CSP-8.3.1 Lupins – Albus No.2 Minimum Receival Standard Farmer Dressed.....76

CSP-10.1.1 Peas – Field No.1 Minimum Receival Standard Farmer Dressed.....77

CSP-10.1.2 Peas – Field No.1 Minimum Export Standard Farmer Dressed..... 78

CSP-10.1.3 Peas – Field No.1 Minimum Export Standard Machine Dressed.....79

CSP-10.2.1 Peas – Field No.2 Minimum Receival Standard Farmer Dressed 80

CSP-10.2.2 Peas – Field No.2 Minimum Export Standard Farmer Dressed.....81

CSP-10.2.3 Peas – Field No.2 Minimum Export Standard Machine Dressed..... 82

CSP-10.3.1 Peas - Field No.3 Minimum Receival Standard Farmer Dressed..... 83

CSP-10.4 Peas – Yellow Split Minimum Export Standard Machine Dressed 84

CSP-11.1 Pigeon Peas - Minimum Receival Standard Farmer Dressed..... 85

CSP-11.2 Pigeon Peas - Minimum Export Standard Farmer Dressed 86

CSP-12.1 Vetch - Minimum Receival Standard Farmer Dressed..... 87

CSP-12.2 Vetch - Minimum Export Standard Machine Dressed.....88

SECTION 4 VARIETY CLASSIFICATION..... 89

SECTION 5 METHODS & PROCEDURES..... 90

5.1 Introduction 90

5.2 Sampling 91

5.3 Moisture Assessment - NIR..... 94

5.4 Defective Grains Assessment – Reference Method 95

5.5 Contaminants & Purity Assessment – Reference Method..... 98

5.6 Sieving Method..... 101

5.7 Sizing Method..... 103

5.8	De-hulling Method.....	104
5.9	Variety Declaration Procedure.....	106
SECTION 6	REFERENCE MATERIALS.....	108

SECTION 1 INTRODUCTION

General

For many years Pulse Australia on an annual basis reviewed, produced and published on behalf of industry all pulse commodity Trading Standards except mung beans (Standards). Grain Trade Australia (GTA) now develops those Standards on behalf of industry through its Trading Standards Committee (Committee).

In order to provide a consistent message to both domestic industry and international buyers, GTA encourages input into the development of these Standards. Additionally, we urge industry to use the Standards contained within this Manual as applicable when buying and trading Australian pulses.

Considerations to the Standards

This section of the Manual relating to pulses has been produced following the annual review by GTA of Standards. There are various sections of this Manual relating to Standards and associated procedures and industry is encouraged to take account of all relevant sections when applying these Standards to pulses bought and traded domestically or internationally.

The Grades referred to in this document are a combination of:

- Grades commonly introduced across the country on an annual basis and are generally the same in each State where pulses are grown or traded. Grades may relate to the receival or outturn (referred to as export) of pulses.
- Grades that may not be introduced every season or only introduced in a regional area. These grades may be created for various reasons including to meet the specific quality requirements of a customer, as a specific varietal segregation or to deal with specific quality issues with harvested grain in a localised area.

Industry should note the list of Grades in this Manual is not exhaustive.

Variations to Standards

Whilst the information in this Manual is current at time of publication, you will need to monitor the GTA Member Updates, the GTA website (www.graintrade.org.au) and other applicable information to ensure that you are aware of the changes to the Standards and the impact on your own trading arrangements.

Varieties

Approved and recommended varieties to be grown and acceptable within each grade or pulse commodity are listed within this document. The process for approval of each variety within a grade is being developed by a group external to GTA.

Variety integrity and correct variety assessment is an integral part of the grain classification and Standards application process. GTA endorses the varietal classifications as listed in this Manual and encourages all industry to follow the approved varietal list as listed in this Manual where relevant.

Changes to varietal classifications may occur at any time during the season following the publication of this Manual. As these changes will not necessarily be included in this Manual industry should implement their own procedures for monitoring the varietal classification process.

Timing of Standards Development

The Standards outlined in this Manual are applicable for the entire season of 2025/26. Standards apply to pulses assessed as per these Standards from 1 August 2025 to 31 July 2026.

SECTION 2 DEFINITIONS

The following definitions have been created to assist in classification of individual pulse grains when using these Standards. The definitions are a general guide, and industry should note that differences to definitions may apply to individual pulse commodities.

The following Defect definitions are to be read in conjunction with the images displayed in the GTA Visual Recognition Standards Guide for Grain Commodity Sampling and Assessment located on the GTA website at www.graintrade.org.au. The images in that document for the applicable pulse commodity display the minimum and/or maximum coverage and attributes of the Defects as defined in these Standards.

Acceptable Sampling Device

Refers to any sampling device that obtains a representative sample to be assessed as per these Standards. Refer to 5.2 for the Sampling Procedures.

Blondes

Applies to Aldinga Red Lentils only.

Parameter	Definition
Cause	A defect causing discolouration of the kernel that may be genetic but potentially influenced by seasonal conditions.
Physical Description	Kernels are not uniformly orange in colour and portions of the kernel appear yellow.
Tolerances to apply	The 'blondes' count is to be included in the total count for Defectives. It is not classified as Poor Colour.
Presence on Seed Coat	n/a
Presence on Kernel	Kernels are not uniformly orange in colour. Kernels appear yellow. Seed coat must be removed to determine the presence on the kernel. Refer also to the GTA Visual Recognition Standards Guide.

Broken & Kibbled

In relation to split product, refers to kernels that fall through the applicable screen. In general, these kernels have a portion missing or are not intact halves.

Broken/Chipped/Loose Seed Coat/Split

Parameter	Definition
Cause	Damage due to poor harvesting and/or handling techniques. Late harvesting may exacerbate this defect.
Physical Description	Breakage, cracking, peeling or splitting of the seed coat or chipping and splitting of the kernel in various forms. Damage to the seed coat may be referred to as loose seed coat or skin damage. Damage to the kernel may be referred to as chipped or scratched.
Presence on Seed Coat	Includes the following: <ul style="list-style-type: none"> ▪ Split Seed Coat - A Split in the seed coat running more than half the entire length or across more than half the entire width on one or both sides. ▪ Skin Damaged - A hole in the seed coat where more than 20% of the seed coat on any one side is missing (Where the entire seed coat is not present, it is often referred to as Missing Seed Coat).

	<ul style="list-style-type: none"> ▪ Loose Seed Coat (Peeling) - Where the seed coat is visibly falling off the kernel to any extent and not adhering tightly to the kernel. ▪ Missing Seed Coat - Where the entire seed coat is missing but the kernel is intact. <p>Refer also to the GTA Visual Recognition Standards Guide.</p>
Presence on Kernel	<p>Any damage to the kernel is classified as defective. Includes the following:</p> <ul style="list-style-type: none"> ▪ Chipped (Scratched) – A part of the kernel is damaged or removed. ▪ A Split where the kernel is separated into two halves. Pieces may be whole or partial. Seed coat may adhere to the kernel pieces (Caps). <p>Refer also to the GTA Visual Recognition Standards Guide.</p>

Bulk Vessel

A sea going vessel used to transport pulses. Pulses are stored loosely in holds without being constrained within a receptacle such as a container or bags. Vessels usually have a number of separate holds or compartments.

Caps

Are any parts of the Seed Coat adhering to Split or Broken seed. Note that the definition refers to “any parts of the Seed Coat”, no matter the size.

Cereal Seeds

In relation to green lentils, red lentils and vetch refers to wheat, durum, barley, oats, sorghum, triticale, cereal rye, maize and rice. Refer also to Foreign Seeds.

Chemicals Not Approved

Refers to the following:

- Chemicals used on the growing crop in the State or Territory where the pulse commodity was grown in contravention of the label.
- Chemicals used on the stored pulse commodity in contravention of the label.
- Chemicals not registered for use on the pulse commodity.
- The pulse commodity containing any artificial colour, pickling compound or marker dye commonly used during crop spraying operations that has stained the pulse commodity.
- The pulse commodity treated with or contaminated by Carbaryl, Organochloride chemicals, or diatomaceous earth.
- Chemical residues in excess of Australian Commonwealth, State or Territory legal limits (see Maximum Residue Limit and National Residue Survey).

For further information on this topic, refer to the document “Australian Grains Industry Post Harvest Chemical Usage Recommendations and Outturn Tolerances 2025/26” - see GTA website <http://www.graintrade.org.au/nwpgp>.

Classification Procedures

Refers to procedures outlined in this document used to sample and assess the quality of pulses tendered for delivery or presented for outturn or export.

Container

A box like receptacle that stores pulses in a sealed environment for transport. Containers are usually approximately 6.1m, 12.2m or 13.7m in length.

Contrasting Colour

Refers to contrasting colour of the seed coat in the same variety. It is usually caused by a recessive gene in that variety. As there is no visible difference on the kernel it is not classified as Defective or Poor Colour. Tolerances apply to the maximum amount permitted in a sample. Refer to the reference Varietal Colour Chart or GTA Visual Recognition Standards Guide for a description of the applicable Contrasting Colours for each variety which depict the minimum and maximum amount of marbling permitted on a seed coat before being classified as Contrasting Colour.

Cotyledon

Refer to the generally recognised term kernel.

Defective Grains

Defective grains refer to a pulse commodity that has been damaged to some degree, due to a range of factors including but not limited to disease, environment, handling, harvest, stress and a weather event. Definitions for Defects apply to the entire seed coat and/or kernel, depending on the defect type and grain type.

Generally, it includes pulses not of the specified variety and seed coats or kernels as follows:

Blondes (Aldinga Red Lentils only)	Insect Damaged
Broken/Chipped/Loose Seed Coat/Split	Severely Damaged
Diseased	Orange Tip (Red Lentils only)
Fungal Affected (e.g., Ascochyta)	Poor Colour
Shrivelled & Wrinkled	Sappy
Green	Sprouted
Hail Damaged	Stained & Weather Damaged
Whole pods containing seed of the pulse being assessed	Seed material of the pulse being assessed passing through a specific sized screen

An individual kernel may only have one defect, being the defect type with the tightest tolerance in the Standard.

a) Sizing of Defect

The size of the quality issue on the seed coat or kernel determines if it is categorised as a defect. Where defined in the tables for each Defect, a specific quality parameter will only be classified as a Defect if its presence exceeds 20% of any one side of the grain even if that Defect is absent or if the coverage is 20% or less on the other side.

As an example, for a Fungal Affected (e.g., Ascochyta) lesion on a Red Lentil, in the following table:

- Highlighted in red is where the grain will be classified as defective due to the large size of the lesion i.e., it exceeds 20%.
- Highlighted in green is where the grain will not be classified as defective due to the small size of the lesion i.e., it is 20% or less.

Relevant Information when classifying grain		Considered a Fungal Affected (e.g., Ascochyta) Lesion?
Lesion Size - Side 1	Lesion Size - Side 2	
5%	15%	N
15%	15%	N
20%	20%	N
25%	5%	Y
30%	10%	Y
10%	30%	Y

b) Defects on Kernels

Where a tolerance is stated for kernels, it is recommended that seed coats be removed to identify the presence of the defect on the kernel. Where the defect is present on the kernel, the grain is classified as Defective except when assessing Poor Colour, where a tolerance may apply. For the definition of Poor Colour, refer to the applicable Standard and grain type.

c) Summary Table for Defect Categories

The following is a table summarising the categories listed within Defective pulses. Note that this table refers to the presence on the entire grain (seed coat or kernel). Variations may exist for some pulses.

Where the statement includes:

- Refer to Definition or Standard – refer to the Definition and relevant description in the applicable commodity Standard.
- Refer to VRSG – refer to the GTA Visual Recognition Standards Guide booklet for further guidance.

Defect Type	Pulse Type	Seed Coat	Kernel
		Classified Defective if greater than	Classified Defective if greater than
General Defects			
Fungal Affected (e.g., Ascochyta)	All pulses except desi chickpeas Desi chickpeas	20% on any one side all pulses except desi chickpeas n/a	Nil (i.e., any presence)
Broken, Chipped, Loose Seed Coat & Split	All pulses	Refer to Definition and VRSG	Nil (i.e., any presence)
Green	Desi Chickpeas	More than a slight greenish tinge	Nil (i.e., any presence)
Poor Colour	All pulses	Refer to Standard & VRSG	Refer to Standard & VRSG
Shrivelled & Wrinkled	All pulses	Distinct Ridge or indentations	Distinct indentation
Severely Damaged	All pulses	Refer to Standard & VRSG	Refer to Standard & VRSG
Stained & Weather Damaged	All pulses	Generally, 20% on any one side. Refer to VRSG	Nil (i.e., any presence)
Varietal Variation Defects			
Blondes	Aldinga Red Lentils	n/a	Nil (i.e., any presence)
Orange Tip	Red Lentils	Refer to VRSG	Nil (i.e., any presence)
Seed Coat - Thin	Desi Chickpeas	Refer to Definition	Refer to Definition
Nil Tolerance Definition Defects			
Hail Damaged	Desi Chickpeas	Nil (i.e., any presence)	Nil (i.e., any presence)
Insect Damaged	All pulses	Nil (i.e., any presence)	Nil (i.e., any presence)
Sappy	All pulses	Nil (i.e., any presence)	Nil (i.e., any presence)
Sprouted	All pulses	Nil (i.e., any presence)	Nil (i.e., any presence)

De-hulling

Is the process of removing the seed coat from the kernel. De-hulling is required to assess the presence of particular defects such as Poor Colour on the kernel.

Export Standard

Refers to the Export Standards outlined in this document. Are Standards that are applied to pulses when sold and transported to overseas markets or outturned on the domestic market.

Farmer Dressed

Refers to pulses that have been harvested and have not subsequently undergone any major cleaning or mechanical screening process to affect their quality. Pulses generally contain some Unmillable Material, Foreign Material and Defective pulses.

Field Fungi

Staining on the seed coat or kernel that may or may not have been caused by the development of fungi during periods of high moisture. Refer to Stained & Weather Damaged for tolerances to apply.

Field Insects

Are live or dead whole insect contaminants that do not cause damage to stored pulses. Tolerances may include dead or live insects depending on the applicable grain type and Standard.

For all Field Insects other than grasshoppers and/or locusts, the definition refers to whole bodies. Body portions or pieces of Field Insects (i.e., dead) are classified as Unmillable Material.

A separate tolerance generally exists for Grasshoppers and Locusts. See also Grasshoppers and Locusts.

For the purposes of these pulse Standards, a Pea Weevil is classified as a Field Insect except for live Pea Weevils. See also Pea Weevil.

Note: There may be variances with the tolerances applied at export by Plant Exports Branch.

Some common Field Insects are listed below:

Common Name	Scientific Name
Ladybirds	<i>Various</i>
Minute Mould Beetles	<i>Corticaria species</i>
Pea Weevil (dead only)	<i>Bruchus pisorum</i>
Sitona Weevil	<i>Sitona species</i>
Desiantha Weevil	<i>Desiantha diversipes</i>
Wood Bugs	<i>Various</i>
All Other Field Insects	

Foreign Material

Refers to Unmillable Material and all vegetable material other than seed material (seed coats or kernels) of the pulse in question being sampled and assessed according to these Standards. This includes:

- Foreign Seeds (Weed seeds), including Small Foreign Seeds
- Cereal seeds (green lentils, red lentils and vetch only)
- Empty seed pods or pieces of seed pods of the pulse being assessed
- Empty seed pods, pieces of seed pods or seed pods containing seeds of all other weed seeds
- Seed attachments of the pulse being assessed
- Unmillable Material (Soil, sand, sticks, Sclerotes etc.)
- Field Insects
- Grasshoppers and Locusts
- Gumnuts

- Ryegrass Ergot
- Snails
- Stored Grain Insects (dead and pieces).

Foreign Material excludes pods containing seeds of the pulse being assessed as these are classified as Defective. Note there may be separate tolerances for parameters listed within Foreign Material.

Foreign Seeds

Are those seeds that are not the pulse being sampled and assessed according to these Standards. Foreign Seeds may be detected both above and below the screen and are to be assessed in the entire sample. Foreign Seeds are included in the assessment of Foreign Material and may also be referred to as Weed Seeds.

Note however, that these weed seed tolerances may differ from those applied in each State and Territory under the respective legislation. All persons trading pulses are advised to refer to the relevant legislation for appropriate Standards to be complied with. GTA takes no responsibility for pulses traded which do not take into account the relevant legislative weed seed standards.

Any seed pods detected must be opened and the seeds counted for inclusion in the tolerances as specified, except where pods have a specified tolerance.

The tolerances listed in each Standard are maximums and refer to the total of all seeds named in each type, except for Type 1 in which the maximum applies on an individual seed basis.

Tolerances for Seed Contaminants apply to whole seeds or their equivalent in pieces per 200 gram, 400 gram or 600 gram sample (above and/or below the screen) of the following species. There may be variations to the following depending on the pulse commodity:

Type 1 Individual Seed Basis

Colocynth (*Citrullus colocynthis*)
 Double Gee / Spiny Emex / Three-Cornered Jack (*Emex australis*)
 Jute (*Corchorus olitorius*)
 New Zealand Spinach (*Tetragonia tetragonioides*)
 Parthenium Weed (*Parthenium hysterophorus*) *
 Poppy (Field) (*Papaver rhoeas*)
 Poppy (Horned) (*Glaucium flavum*)
 Poppy (Long Head) (*Papaver dubium*)
 Poppy (Mexican) (*Argemone mexicana*)
 Poppy (Wild) (*Papaver hybridum*)

* QLD only. A NIL tolerance applies in NSW/VIC/SA

Type 2

Castor Oil Plant (*Ricinus communis*)
 Coriander (*Coriandrum sativum*)
 Crow Garlic / Wild Garlic (*Allium vineale*)
 Darling Pea (*Swainsona spp*)
 Opium Poppy (*Papaver somniferum*)
 Peanut seeds and pods (*Arachis hypogaea*)
 Ragweed (*Ambrosia spp*)
 Rattlepods (*Crotalaria spp*)
 Starburr (*Acanthospermum hispidum*)
 St. John's Wort (*Hypericum perforatum*)

Type 3 (a)

Bathurst Burr (*Xanthium spinosum*)
 Branched Broomrape (*Orobanche ramosa*)
 Bulls Head / Caltrop / Cats Head (*Tribulus terrestris*)
 Cape Tulip (*Homeria spp*)
 Cottonseed (*Gossypium spp*)
 Dodder (*Cuscuta spp*)

Noogoora Burr (*Xanthium pungens*)
Thornapple (*Datura spp*)

Type 3 (b)

Vetch (Commercial) (*Vicia spp*) *
Vetch (Tare) (*Vicia sativa*) *

* Nil tolerance applies to vetch of any type in Red Lentils (whole or split) to Saudi Arabia

Type 3 (c)

Heliotrope (Blue) (*Heliotropium amplexicaule*)
Heliotrope (Common) (*Heliotropium europaeum*)

Type 4

Bindweed (Field) (*Convolvulus arvensis*)
Cutleaf Mignonette (*Reseda lutea*)
Darnel (Drake Seed) (*Lolium temulentum*)
Hexham Scent / Melilot (King Island) (*Melilotus indicus*) - only acceptable if no tainting odour is present
Hoary Cress (*Cardaria draba*)
Mintweed (*Salvia reflexa*)
Nightshades (*Solanum spp*)
Paddy Melon (*Cucumis myriocarpus*)
Skeleton Weed (*Chondrilla juncea*)
Variegated Thistle (*Silybum marianum*)

Type 5

Knapweed (Creeping / Russian) (*Acroptilon repens*)
Patterson's Curse / Salvation Jane (*Echium plantagineum*)
Sesbania Pea (*Sesbania cannabina*)

Type 6

Clover (Pods) (*Trifolium spp*)
Colombus Grass (*Sorghum alnum*)
Johnson Grass (*Sorghum halepense*)
Lucerne (Pods) (*Medicago spp*)
Marshmallow (Pods) (*Malva parviflora*)
Medic (Pods) (*Medicago spp*)
Muskweed (Pods) (*Myagrurn perfoliatum*)
Saffron Thistle (*Carthamus lanatus*)
Sow Thistle (Heads) (*Sonchus spp*)
Trefoil (Pods) (*Medicago spp*)
Wild Radish (Pods) (*Raphanus raphanistrum*)

Pods refers to whole pods or part thereof.

Type 7 (a)

Adzuki Beans (*Vigna angularis*)
Broad beans (*Vicia faba*)
Chickpeas (Desi/Kabuli) (*Cicer arietinum*)
Corn (Maize) (*Zea mays*)
Cowpea (*Vigna unguiculata*)
Faba Beans (*Vicia faba*)
Lentils (Green/Red) (*Len culinaris*)
Lupin (Albus/Angustifolius) (*Lupinus spp*)
Mung beans (*Vigna radiata*)
Onion Weed Pods (*Asphodelus fistulosus*) regardless of size
Peas (Field) (*Pisum sativum*)
Pigeon Peas (*Cajanus cajan*)
Soybean (*Glycine max*)

Excludes the pulse being sampled.

Includes any other seeds or pods greater than 5mm in diameter not listed in the Standards.

Type 7 (b)

Barley (2 row) (*Hordeum distichon*)
 Barley (6 row) (*Hordeum vulgare*)
 Bindweed (Australian) (*Convolvulus erubescens*)
 Bindweed (Black) (*Polygonum convolvulus*)
 Brome Grass (*Bromus spp*)
 Carrot Weed (*Daucus carota*)
 Durum (*Triticum durum*)
 Oats (Black/Wild) (*Avena fatua*) *
 Oats (Common) (*Avena sativa*) *
 Oats (Sand) (*Avena strigose*) *
 Red / Spring Feed Wheats (Various)
 Rice (*Oryza sativa*)
 Rye (Cereal) (*Secale cereale*)
 Ryegrass on Stalk (*Lolium spp*)
 Sorghum (Forage) (*Sorghum bicolor*)
 Sorghum (Grain) (*Sorghum bicolor*)
 Triticale (*Triticosecale spp*)
 Turnip Weed Pods (irrespective of size) (*Rapistrum rugosum*)
 Wheat (*Triticum aestivum*)

Other than Small Foreign Seeds, includes any other seed contaminant not specified in Standards.

Note: A separate tolerance exists for Cereal Seeds in Green Lentils, Red Lentils and Vetch only. Refer definition of Cereal Seeds.

* Individual seeds in cluster to be counted.

Type 7 (c)

Safflower (*Carthamus tinctorius*)
 Sunflower (*Helianthus annuus*)

Type 8

Bellvine (*Ipomoea plebera*)

Fungal Affected (e.g., Ascochyta)

Parameter	Definition
Cause	Is a fungal disease that attacks plant foliage and seed pods
Physical Description	Lesions are generally visible to the naked eye. The lesion generally appears intense dark brown to black. It is commonly oval to circular and localised in nature but may vary in shape. The lesion may be similar in colour to Severely Damaged or Stained & Weather Damaged. The lesion may also be associated with the presence of fungal growth of various colours. A lesion may appear on one or both sides of the seed coat or kernel.
Presence on Seed Coat	Lesion greater than 20% coverage on any one side of the seed coat for all pulses except desi chickpeas. For all grains, the visual presence of any level of fungal growth associated with mould is considered defective. Refer also to the GTA Visual Recognition Standards Guide. Refer also to Stained & Weather Damaged. For Desi Chickpeas, any lesion of any size is permitted and not classified as Fungal Affected provided it is not also present on the kernel. For Desi Chickpeas, if the Fungal Affected seed coat lesion is >20% but does not penetrate to the kernel (and

	thus fall under the Fungal Affected definition), then the grain is classified as Stained & Weather Damaged and is classified as defective.
Presence on Kernel	Any lesion of any size present on the kernel is classified as defective.

Good Colour

Are those seed coats and kernels practically free from discolouration and have the uniform natural colour and appearance characteristic of the predominating pulse type within the sample of the pulse being assessed. The colour and thus definition may vary by pulse. Note that dark colours such as black may be excluded as the predominating colour depending on the pulse type. Refer to each Standard and the GTA Visual Recognition Standards Guide for details.

A thin, transparent seed coat may allow the orange kernel to show through in red lentil varieties such as Aldinga and Nugget, hence the seed appears 'orange'. This is deemed acceptable and not Poor Colour. For all commodities, seeds that have partial transparency, but an acceptable kernel colour are deemed acceptable and of Good Colour. Refer also to Contrasting Colour.

Grasshoppers and Locusts

For grasshoppers and/or locusts, six legs, three body parts or two wings or part thereof, constitutes one entire insect respectively. More than one of the same body part constitutes greater than one insect.

Tolerances may apply to live and/or dead, depending on the commodity and Standard.

Grasshoppers and Locusts may have a separate tolerance or be included in the tolerance for Field Insects depending on the commodity and Standard. Refer also to Field Insects.

Green Grains

Variations to the following definitions and tolerances may apply depending on the commodity and Standard.

Parameter	Definition
Cause	Premature ripening of the grain.
Physical Description	Seed coats or kernels appear green. Where any greenish tinge is present on the seed coat, it is recommended the kernel also be inspected.
Presence on Seed Coat	For most commodities, more than a slight greenish tinge must be present to be classified as defective. Refer also to the GTA Visual Recognition Standards Guide.
Presence on Kernel	For most commodities, any damage to the kernel is classified as defective. Refer also to the GTA Visual Recognition Standards Guide.

Grower Load Composite

A sample representing the entire load tendered for delivery. Compiled by obtaining individual probe samples of the individual load (container, truck etc.) based on the tonnage each represents and combining these samples to form one sample. Refer to 5.2 for the Sampling Procedure.

Gumnuts

A gumnut is defined as the woody fruit of trees of the genus Eucalyptus. The tolerance refers to whole or pieces of any size and maturity level.

Hail Damaged – Desi Chickpeas

Parameter	Definition
Cause	Damage by hail.
Physical Description	Damage to the seed coat or kernel. Damage to the seed coat can appear as bruising (darkening) or in more severe cases splitting of the seed coat. This may cause discolouration and damage to the kernel. Damage to the kernel can vary from bruising (darkening) to physical damage such as crushing of the entire kernel.
Presence on Seed Coat	Any damage to the seed coat is classified as defective. Refer also to the GTA Visual Recognition Standards Guide.
Presence on Kernel	Any damage to the kernel is classified as defective.

Insect Damaged

Parameter	Definition
Cause	Damage due to any insect such as Pea Weevil, Etiella grub and Heliothis eating the seed coat or more commonly, the kernel.
Physical Description	The seed coat and kernel have a chewed appearance. Kernels may contain holes as a result of insects boring through the kernel. Mechanical damage resulting in Broken or Split grains is not included in this definition.
Presence on Seed Coat	Any damage to the seed coat is classified as defective. Damage generally also occurs under the seed coat and is obvious on the kernel. Includes any number of insect eggs of all Stored Grain Insect species on the Seed Coat surface, whether those eggs may be live or dead. Refer also to the GTA Visual Recognition Standards Guide.
Presence on Kernel	Any damage to the kernel is classified as defective. Refer also to the GTA Visual Recognition Standards Guide.

Kernel

Refers to the inner part of a pulse that is contained under the seed coat. It may also be referred to as a Cotyledon. Note that for most commodities and in most Standards, any damage to the kernel results in that grain being classified as Defective except for Poor Colour.

Load

A road bulk unit tendered for delivery.

Loose Seed Coat

Refer to Broken/Chipped/Loose Seed Coat/Split.

Machine Dressed

Refers to pulses that have undergone a significant quality transition via a mechanical operation such as cleaning to remove Foreign Material, Foreign Seeds or Defective pulses.

Manganese Deficiency

In relation to *Angustifolius* lupins refers to splitting of the seed coat to expose the kernel. It is not considered a defect provided no damage to the exposed kernel has occurred. Refer to the GTA Visual Recognition Standards Guide for lupins.

Maximum Residue Limit (MRL)

MRLs are the maximum amount of a chemical residue or its metabolite that is legally permitted on or in agricultural commodity. The Australian Pesticides and Veterinary Medicines Authority (APVMA) sets MRLs. These MRLs are set at levels which are not likely to be exceeded if the agricultural or veterinary chemicals are used in accordance with approved label instructions and can be found on the ComLaw website at the following link <https://www.legislation.gov.au/F2023L01350/latest/versions>

Australian MRLs may differ significantly from those prescribed by foreign countries and the International Codex Alimentarius Commission. Consequently, grain exporters must be aware of MRLs of importing countries and which countries accept Codex MRLs. Foreign country MRLs may be accessed directly from foreign government websites or the NRS grains database at <https://www.agriculture.gov.au/agriculture-land/farm-food-drought/food/nrs/databases>

Missing Seed Coat

Refers to those grains where the Seed Coat is partially or more usually, entirely missing. It may also be referred to as “Seed-Coatless”. Refer to Broken/Chipped/Loose Seed Coat/Split. Refer also to each Standard and the GTA Visual Recognition Standards Guide for details.

Moisture Content

The amount of water measured in a sample of pulses, being assessed according to these Standards.

National Residue Survey

The National Residue Survey (NRS) gathers information and supplies chemical residue results on domestic and export grain commodities. Australian grain is of a high quality with respect to residues and contaminants. It is recommended all grain exporters, container packers, bulk export terminal operators, Bulk Handling Companies and processors actively participate in the NRS grains residue monitoring program. Contravention of an overseas MRL may cause the rejection of cargoes resulting in severe financial cost being incurred and potentially jeopardising Australian grain into that market. Information about the NRS is located: <https://www.agriculture.gov.au/agriculture-land/farm-food-drought/food/nrs>

Nil

Means a level of zero in a 200 gram, 400 gram or 600 gram sample representative of the entire load. Nil means not detected anywhere in the load or at any stage of the receipt or unloading process.

Non-vegetable Matter

Is a part of Unmillable Material. It refers to all non-organic material, including Soil and Stones. Different tolerances may apply depending on the material and pulse type.

Objectionable Material

Refers to any objectionable foreign material that may or may not be otherwise stated in these Standards which has the ability to degrade the hygiene of the pulse, become a food safety issue of concern or has a commercially unacceptable odour (refer Objectionable Odour).

This includes but is not limited to the following:

- Animal excreta.
- Rodents, either alive or dead.
- Crushed insect bodies or parts that adhere to the grain causing clumping of the grain that may or may not cause an odour.
- Live Stored Grain Insects.
- Any chemical not registered for use on pulses, in excess of legal tolerances or chemicals in excess of the MRL.
- Pickling compounds or Artificial colour / seed dressings or any fungicide added to the pulse as a seed dressing.
- Any tainting agents and / or other contaminants imparting an odour not normally associated with that particular pulse.

- A commercially objectionable odour and/or an odour not normally associated with the pulse in question. Odour may be caused by various means which may or may not be discernible in the sample being assessed.
- Toxic and / or noxious weed seeds which are prohibited by State laws against inclusion in stockfeed.
- Ergot of any commodity except Ryegrass Ergot (for which a tolerance may apply).
- Any other commercially unacceptable contaminant such as glass, metal, fertiliser, concrete or other contaminants greater than the tolerance or specifications as allowed in the Standards.

As many of these parameters such as chemicals are not able to be assessed on site prior to delivery of the pulse, it is the responsibility of the grower or deliverer of the pulse to ensure compliance with any regulations or Standards. It is recommended that a declaration be obtained by the Storage Provider regarding the chemical status of the pulse tendered for delivery.

Objectionable Odour

In the context of these Standards is a commercially objectionable odour and/or an odour not normally associated with the pulse in question. The Objectionable Odour may be caused by various means which may or may not be discernible in the sample being assessed. A nil tolerance applies.

Orange Tip

Applies to Red Lentils only.

Parameter	Definition
Cause	A defect causing discolouration of the seed coat caused by seasonal conditions. It may or may not be a pre-cursor to sprouting.
Physical Description	Seed coats have varying degrees of orange colour, generally in the germ area.
Tolerances to apply	Orange tip is to be included in the count for Total Defectives. It is not classified as Poor Colour.
Presence on Seed Coat	Seed coats appear orange in colour. Refer also to the GTA Visual Recognition Standards Guide.
Presence on Kernel	n/a

Outturn

Process of loading the pulse from a storage unit into a transport unit, for eventual delivery to a domestic or international customer. Standards as listed in this document may apply.

Pea Weevil

Refers to any stage in the life cycle of insects of the species *Bruchus pisorum*. For the purposes of these pulse Standards, a Pea Weevil (dead only) is classified as a Field Insect. See also Field Insects.

<u>Parameter</u>	<u>Classified as</u>
Pea Weevil - live	Objectionable Material (Nil tolerance)
Pea Weevil - dead, entire	Field Insects
Pea Weevil – dead, pieces	Unmillable Material

A NIL tolerance applies to live Pea Weevil as they are classified as Objectionable Material.

Physical Characteristics

Is a general description of pulses. Usually describes the general appearance and overall condition relative to a particular variety of the relevant pulse type.

Phomopsis

Is a fungal disease that causes various agronomic and quality issues in pulses such as lupins. Refer to each Standard and the GTA Visual Recognition Standards Guide for details.

Pickling Compounds or Artificial Colour

Grains contaminated in this way may be identified by an unnatural surface colour and/or a colour that rubs off. Any grains that are artificially coloured regardless of intensity, coverage or colour are defective.

Plant Exports Branch

The Plant Export Operations Branch facilitates trade and market access through the development of technical operational phytosanitary export inspection and certification policies. The branch does this by developing and implementing policies and guidance materials that ensure Australian horticulture, grain and other plant-based exports meet Australia's export legislation, international obligations and importing country conditions. The branch has a key role in advancing Australia's market access priorities by actively engaging in overseas market access negotiations. The branch works with all participants in the export pathway to continuously improve the export framework and ensure expert advice and support is available to the horticulture and grain export industries. In addition, the branch administers the development and training of authorised officers to conduct inspections.

Poor Colour

Parameter	Definition
Cause	Rapid, premature ripening. Discolouration may also arise through weather conditions, disease or during the storage period
Physical Description	Poor Colour seed coats or kernels are not considered good colour. Seed coats and kernels vary from white to dark brown/black, and green, depending on the pulse type. Refer to each Standard and the GTA Visual Recognition Standards Guide for further information. Seed coats and kernels may be similar in appearance to various other defects such as Severely Damaged. Does not include Contrasting Colour. Refer also to the definition for Contrasting Colour.
Presence on Seed Coat	Refer to each Standard and the GTA Visual Recognition Standards Guide for further information.
Presence on Kernel	Refer to each Standard and the GTA Visual Recognition Standards Guide for further information. Seed coat must be removed to determine the presence on the kernel.
Variety Specific Definitions not elsewhere listed	
Faba Beans	Grey seed coats due to staining from the seed pod lining. Refer to the GTA Visual Recognition Standards Guide.
Faba Beans	Pea Seed Borne Mosaic Virus – Staining on the seed coat caused by the Pea Seed Borne Mosaic Virus. Refer to the GTA Visual Recognition Standards Guide.
Desi Chickpeas	Seed coat – is included under Total Defects, generally described as Stained/Weather Damaged. Kernel – a separate tolerance exists under the Total Defects category. Refer to the GTA Visual Recognition Standards Guide.
Kabuli Chickpeas	Seed Coat – where seed coats are “bright white / bleached”, included in Poor Colour.

Predominating Class

Is used in reference to the determination of Poor Colour. Refers to the overall colour of a sample where grains in the greatest quantity within the sample are considered to be Good Colour. The Predominating Class specifically excludes grains not considered to be Good Colour. Refer also to Contrasting Colour and the applicable commodity Standard for further details.

Purity

The amount of seed material of the particular pulse in question in the sample. Purity only includes the seed coat and kernel whether intact or defective. It excludes all other plant material of the pulse in question. Purity is generally the opposite of Foreign Material.

Quarantine Requirements

Are those parameters that are mandated by law by an importing country Government Quarantine Authority that must be met in order to permit entry of the particular pulse. Quarantine requirements are also mandated in Australian Export Legislation. On export of pulses from Australia, these quarantine regulations are enforced by Plant Exports Branch. Refer to Plant Exports Branch.

Receival Standard

Refers to the Receival Standards as outlined in this document. Receival Standards apply to the purchase of pulses from a grower or through the Trade.

Representative Sample

A sub-sample of a parcel of pulses used for assessment purposes, which is representative of the entire pulse parcel.

Ryegrass Ergot

Is a contaminant resulting from the infection of ryegrass kernels by the fungus *Claviceps purpurea*.

The tolerance generally applies to the length in centimetres of all pieces lined end to end.

Sappy

Parameter	Definition
Cause	Are those grains that have been harvested before maturity.
Physical Description	Grains are generally soft when pressed.
Presence on Seed Coat	Any level of sappiness is classified as defective.
Presence on Kernel	Any level of sappiness is classified as defective.

Sclerotes

Sclerotes are fungal disease survival structures arising from fungal infection. Refer also to Unmillable Material.

Seed Coat

The outer surface of many pulses that envelopes the kernel. Its function is to protect the kernel from splitting or being damaged. The seed coat is often paper-thin.

Seed Material

Whole seeds, pieces of seed coats and kernels of the pulse being assessed.

Seed Pod

The protective enclosure, shell, or case surrounding a seed or a number of seeds.

Severely Damaged

Parameter	Definition
Cause	<p>A range of causes including:</p> <p>Exposure to bacteria or fungi in the field or in storage.</p> <p>Exposure to severe heat during storage. Heating occurs via mould damage or incorrect drying of high moisture grain.</p> <p>Heat, subsequent mould attack and high moisture conditions may lead to adherence of foreign material or joining of mouldy grains.</p> <p>Various other factors.</p>
Physical Description	<p>Damage to the grain causing it to become severely discoloured. A grain exhibits one or more of the following characteristics:</p> <p>Heat Damaged or Burnt:</p> <p>Heat Damaged or Burnt refers to those grains where the seed coat or kernel has become severely discoloured. Affected grains appear dark brown or in severe cases, blackened.</p> <p>Mould:</p> <p>Affected grains appear discoloured and visibly affected by mould.</p> <p>Other Serious Visual Defects:</p> <p>Refers to those grains that have become significantly discoloured and / or have a serious visual defect that is not otherwise listed in these Standards.</p> <p>This definition does not include Fungal Affected (e.g., Ascochyta) lesions or known Diseases such as Botrytis Grey Mould. An Objectionable Odour must not be detected.</p>
Presence on Seed Coat	Damage to the seed coat is classified as defective. Refer to the GTA Visual Recognition Standards Guide for the minimum damage required.
Presence on Kernel	Any damage to the kernel is classified as defective. Refer to the GTA Visual Recognition Standards Guide.

Shrivelled & Wrinkled

Parameter	Definition
Cause	Damage has occurred during the maturation phase due to some form of environmental or agronomic stress such as frost.
Physical Description	Visible damage to the seed coat or size and shape of grain whereby the grains are severely distorted and/or shrunken. Seed coats may show a level of discolouration depending on the extent of damage. Grains are often smaller than the majority in the sample. For some commodities, these small grains may fall through the screen and be classified as Defective.
Presence on Seed Coat	A distinct ridge (often described as mountains and valleys) on the seed coat must be present to be classified as Shrivelled & Wrinkled. Ridges may be described as coarse waves rather than soft waves. Seed coats may be wrinkled or dimpled and distinctly indented into the kernel. Seed coats with a slight degree of indentations are not included in this definition. Refer also to the GTA Visual Recognition Standards Guide.
Presence on Kernel	A distinct indentation on the kernel.

Sizing

Refers to the procedure used to determine the amount of pulse material retained above the specified screen, following removal of all Foreign Material. Sometimes referred to as Retention.

Skinless

Refers to those kernels with a Missing Seed Coat. Refer to Broken / Chipped / Loose Seed Coat / Split.

Small Foreign Seeds

These are foreign seeds that are small in size. They are included in the Foreign Material fraction and generally fall below the screen during the screening process, where a screen is used, except for those foreign seeds specifically listed in the Foreign Seeds Type categories. A list of the more common Small Foreign Seeds is below. Note that not all Small Foreign Seeds may be listed.

Common Name	Scientific Name
Amsinckia	<i>Amsinckia spp</i>
Australian Phalaris	<i>Phalaris aquatica</i>
Bladder Soapwort	<i>Vaccaria hispanica</i>
Broomrape (Clover/Common)	<i>Orobanche minor</i>
Burrweed (Yellow)	<i>Amsinckia spp</i>
Canary Grass (Australian)	<i>Phalaris aquatica</i>
Canary Grass (Lesser)	<i>Phalaris minor</i>
Canary Grass (Wild)	<i>Phalaris canariensis</i>
Canola	<i>Brassica rapa</i>
Celery (Slender)	<i>Apium leptophyllum</i>
Charlock	<i>Sinapis arvensis</i>
Clover (Ball)	<i>Trifolium glomeratum</i>
Cockspur (Maltese)	<i>Centaurea melitensis</i>
Dock	<i>Rumex spp</i>
Evening Primrose	<i>Oenothera stricta</i>
Fat Hen	<i>Chenopodium album</i>
Fescue	<i>Festuca spp</i>
Hares Ear	<i>Conringia orientalis</i>
Hedge Mustard	<i>Sisymbrium officinale</i>
Horehound	<i>Marrubium vulgare</i>
Juncea Canola	<i>Brassica juncea</i>
Knotweed	<i>Polygonum aviculare</i>
Lettuce	<i>Lactuca spp</i>
Lucerne (Seeds)	<i>Medicago sativa</i>
Maltese Cockspur	<i>Centaurea melitensis</i>
Marshmallow (Seeds)	<i>Malva palviflora</i>
Medic (Seeds)	<i>Medicago spp</i>
Muskweed (Seeds)	<i>Myagrurn perfoliatum</i>
Mustard (Wild)	<i>Sisymbrium spp</i>
Mustard (Indian Hedge)	<i>Sisymbrium orientale</i>
Paradoxa Grass (Seeds)	<i>Phalaris paradoxa</i>

Common Name	Scientific Name
Peppercress	<i>Lepidium spp</i>
Phalaris (Australian) (Seeds)	<i>Phalaris aquatica</i>
Radish (Wild) (Seeds)	<i>Raphanus raphanistrum</i>
Rapeseed	<i>Brassica rapa</i>
Ryegrass	<i>Lolium spp</i>
Sage (Wild)	<i>Salvia verbenaca</i>
Saltbush	<i>Atriplex spp</i>
Sorrel	<i>Rumex acetosella</i>
Sowthistle	<i>Sonchus spp</i>
Thistle (Milk) (Seeds)	<i>Sonchus oleraceus</i>
Turnip (Mediterranean)	<i>Brassica tournefortii</i>
Turnip (Wild)	<i>Brassica rapa</i>
Urochloa Grass	<i>Urochloa panicoides</i>
Verbena	<i>Verbena spp</i>
Wireweed	<i>Polygonum aviculare</i>

Snails

Refers to whole or substantially whole (more than half) empty snail shells, bodies or bodies with shells, irrespective of species. Tolerances generally apply to live and dead snails. Pieces of material not defined as a Snail (i.e., smashed snail shells that remain in the sample after cleaning) are classified as Unmillable Material.

Soil

Is generally regarded as unconsolidated mineral or organic material. Soil comprises clumps or grains of earth and grains of sand. Is included in the definition of Unmillable Material, however a separate tolerance for Soil generally applies.

Recognising the inevitability of a small level of Soil contamination at harvest, and in order to provide a practical standard that recognises both the difficulty for delivery of Farmer Dressed pulses completely free of Soil, and the requirement of most importing countries for zero tolerance of Soil in imported product:

- The Receival Standards for almost all Farmer Dressed pulses have been set at a maximum of 0.5% total Unmillable Material by weight, which includes a maximum of 0.3% of Soil by weight.
- For most Machine Dressed product, the Export Standards have been set at a maximum of 0.1% of Unmillable Material by weight, which includes Soil.

Note there is no size limit on the definition of Soil.

These Pulse Standards have been set on the basis of past experience, which has shown that at the levels set, any Soil present at receival is likely to dissipate through the normal handling and/or settling processes to the point of being undetectable.

All industry participants should, however, be aware that most importing countries prohibit any Soil contamination whatsoever, and it is the responsibility of all individuals involved in the trade of pulses to confirm for themselves prior to shipment, that Soil levels in any consignment conform with the specifications of their contract and/or the legal requirements of the importing country and will pass the Plant Exports Branch inspection process.

For further guidance on importing country quarantine requirements, refer to the Department of Agriculture, Fisheries and Forestry Micor database located at <https://micor.agriculture.gov.au/Plants/Pages/default.aspx>

Speckling

Speckling arises from a genetic stress during maturation or may be a genetic characteristic of that variety. It does not refer to black grains for which a tolerance may apply in the Poor Colour category.

Speckling may appear as a few to many small spots of varying colours. For all commodities, Speckling is not considered a defect when it appears to any extent on the Seed Coat. If Speckling is on the Kernel, it is considered a defect.

Red Lentils - Spots may be few or cover a significant portion of the seed coat or kernel. Refer also to Contrasting Colour in the GTA Visual Recognition Standards Guide.

Desi Chickpeas - commonly appears as small spots of any colour on the seed coat or kernel. Spots may be few or cover a significant portion of the seed coat. Refer to the GTA Visual Recognition Standards Guide.

Split

Refer to Broken/Chipped/Loose Seed Coat/Split.

Sprouted

Parameter	Definition
Cause	Damage due to wet weather conditions during maturation. Also occurs through moisture ingress when in storage.
Physical Description	The seed coat has split, and the primary root has emerged. This includes early and any further advanced stage of growth of the primary root. Includes grains where the primary root has been knocked off during the harvesting or handling process.
Presence on Seed Coat	Any visual presence of the primary root through the seed coat is classified as defective. Refer also to the GTA Visual Recognition Standards Guide. Where the seed coat has split but the primary root has not emerged, the grain is to be classified as per the definition of Broken, Chipped, Loose Seed Coat and Split.
Presence on Kernel	Any visual presence of the primary root through the seed coat is classified as defective. Kernels may also be soft to the touch. Refer also to the GTA Visual Recognition Standards Guide.

Stained & Weather Damaged

Parameter	Definition
Cause	Damage has occurred during the maturation phase due to some form of disease, weather event, adherence of Foreign Material or stress prior to harvest.
Physical Description	A general term used to describe visible damage to the seed coat or kernel that may or may not otherwise be defined or be distinguishable from other defects in these Standards. Seed coats and kernels may be discoloured or altered in size or shape. Stained & Weather Damaged is included in Poor Colour, and may also lead to a Loose Seed Coat, Shrivelled and Wrinkled.

Presence on Seed Coat	<p>Discolouration of the seed coat of various shapes and shades. Generally, is a dark brown to black colour depending on the pulse type. May be on one or more sides. Depending on the intensity and the pulse type, generally must be greater than 20% of the surface area on any one side of the seed coat. Refer also to the GTA Visual Recognition Standards Guide.</p> <p>May also appear as some form of light Staining due to minor levels of adherence of Foreign Material (i.e., dust, soil).</p> <p>For Desi Chickpeas, if the Fungal affected (e.g., Ascochyta) seed coat lesion is >20% but does not penetrate to the kernel (and thus fall under the Fungal Affected definition), then the grain is classified as Stained & Weather Damaged and is classified as defective.</p>
Presence on Kernel	Any damage to the kernel is classified as defective.

Sticks

Refers to ligneous material of any size. Sticks are included in Foreign Material.

Stones

Refers to a lump or mass of hard consolidated mineral matter of any size. Stones are included in Unmillable Material.

Stored Grain Insects

Are insect contaminants that generally cause damage to the stored pulse. There is a nil tolerance for live Stored Grain Insects at any life stage of development (i.e., eggs, pupae, larvae and adults) in all Standards. This nil tolerance applies to all Stored Grain Insects specified in the Standards, and those considered as a Stored Grain Insect at export, as listed in the Export Control Act 2021 as implemented by the Department of Agriculture, Fisheries and Forestry.

Dead or pieces of Stored Grain Insects are included in Unmillable Material.

Commonly found Stored Grain Insects include the following:

Common Name	Scientific Name
Angoumois Grain Moth	<i>Sitotroga cerealella</i>
Confused Flour Beetle	<i>Tribolium confusum</i>
Cowpea Bruchid	<i>Callosobruchus maculatus</i>
Flat Grain Beetle	<i>Cryptolestes spp.</i>
Granary Weevil	<i>Sitophilus granaries</i>
Indian Meal Moth	<i>Plodia interpunctella</i>
Lesser Grain Borer	<i>Rhyzopertha dominica</i>
Maize Weevil	<i>Sitophilus zeamais</i>
Psocids/Book Lice	<i>Psocoptera spp.</i>
Rice Weevil	<i>Sitophilus oryzae</i>
Rust-red Flour Beetle	<i>Tribolium castaneum</i>
Saw Toothed Grain Beetle	<i>Oryzaephilus surinamensis</i>
Tropical Warehouse Moth	<i>Ephestia cautella</i>
Warehouse Beetle	<i>Trogoderma variable</i>

For Stored Grain Insect eggs on the surface of a pulse Seed Coat, refer to Insect Damaged.

Thin Seed Coat

Applies to Desi Chickpeas only.

Parameter	Definition
Cause	A defect causing transparency or removal of the Seed Coat. It may or may not be involved with removal of the radicle. It may be a pre-cursor to sprouting.
Physical Description	The Seed Coat is "thin", resulting in the kernel being observed below the Seed Coat. Not Contrasting Colour.
Tolerances to apply	Grain to be classified as Sound or Broken / Chipped / Loose Seed Coat / Split as noted below.
Presence on Seed Coat	Desi radicle is intact, the Seed Coat also is intact but a small section on the radicle is "thin", resulting in the kernel being observed below the Seed Coat. Classify grain as sound.
Presence on Kernel	Where the Seed Coat has been removed and the Desi radicle is not intact (i.e., broken to any extent or removed), classify as Broken / Chipped / Loose Seed Coat / Split

Tiger Striping

In relation to Desi Chickpeas, Tiger Striping is typically due to a period of high heat stress during grain maturation or is a varietal characteristic. Tiger Striping may appear as dark coloured lines of striping or sometimes extending down the seed coat as a mass of colour. Tiger Striping is not considered a defect when it appears to any extent on the Seed Coat. If Tiger Striping is on the Kernel, it is considered a defect Refer to the GTA Visual Recognition Standards Guide.

Taint

Arises from contaminants imparting any smell or taint to the pulse. Includes but is not limited to plant parts and seeds of *Eucalyptus spp* except Gumnuts for which a separate tolerance may apply. Refer also to Objectionable Odour.

Unmillable Material

Includes Soil, Sand, Stones, pieces of Snail, pieces of Stored Grain Insects, pieces of Field Insects (except Grasshoppers and Locusts), Sclerotes and other non-vegetable matter. Refer also to Objectionable Material.

Varietal Colour Charts

This parameter mainly relates to red lentils. Using a physical description and photographs, these charts depict the normal colour and colour range of each variety where a tolerance for Contrasting Colour applies. The charts also indicate the colours that are considered as Contrasting Colour for which a tolerance may exist in the Standards. There is a requirement for a chart to be produced for all varieties where a tolerance for Contrasting Colour exists. Refer also to the GTA Visual Recognition Standards Guide.

Varietal Masterlist

This list designates the varietal group into which each variety may be assigned. For all commodities except Mung Beans, there is generally no restriction on the grade each variety may be received into.

Variety

This is the next lowest level taxonomic rank of a plant below that of the term "species". Differing varieties have differing genetic compositions which may endow them with differing agronomic characteristics, and/or differing end product quality characteristics.

Visual Recognition Standards Guide

The Visual Recognition Standards Guide (VRSG) contains a range of photographs and illustrations to supplement the Standards as outlined in this booklet for particular pulse commodities. The most recent VRSG for various pulses was released by Grain Trade Australia in August 2025. It can be obtained at the GTA website www.graintrade.org.au

To assess Broad Beans, refer to the Faba Bean section of the VRSG.

Whole

Refers to an individual pulse where the seed coat and kernel are entirely 100% present. If part of the seed coat or kernel has been removed due to poor harvesting and handling techniques or through other means, then these are not considered Whole.

Whole Pods

Refers to a fully intact protective enclosure, shell, or case surrounding a seed. Whole pods may contain several seeds.

SECTION 3 GRAIN QUALITY STANDARDS

The following tables represent the grades of various pulses as defined in this Manual.

Standards exist for various commodities in various forms, including:

- At Receival – Farmer Dressed
- At Export – Farmer Dressed or Machine Dressed
- Other Processed commodities

All Standards are referred to by a CSP number, unique to each Standard.

To fully understand and accurately implement the pulse Quality Standards, reference should be made to other relevant sections in this Pulse Manual including:

- Definitions
- Varietal Master List
- Methods & Procedures
- Reference materials such as the Visual Recognition Standards Guide

Other sections of the GTA Standards Manual should also be perused for general guidance on activities associated with implementation of these Standards.

As stated previously, the following Standards are applicable at the time of publishing of this Manual. Variations and new Grades may exist, and industry is encouraged to keep updated with changes via reviewing the GTA website and other relevant industry information sources.

Industry should note that varieties listed within these Standards may be subject to Plant Breeder's Rights.

CSP-1.1 Adzuki Beans Minimum Receival Farmer Dressed

(Grain Trade Australia) ADZUKI BEANS STANDARDS 2025/2026	
Commodity: Adzuki Beans	Season: 2025/2026
Grade: Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP 1.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS The Adzuki Beans shall be sound, dry, fresh and have true varietal colour and characteristics for "Bloodwood" and "Erimo".	
Purity Minimum (% by weight). Whole Adzuki Beans, defective Adzuki Beans and Adzuki Beans seed coats.	99
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Adzuki Beans not of the specified variety and Adzuki Beans that are broken, chipped, damaged, diseased, frost damaged, insect damaged, loose seed coat, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed, and all Adzuki Bean seed material falling below the 4.33mm round hole screen.	5
of which, Split/Broken Maximum (% by weight). Adzuki Beans that are not whole.	3
of which, Damaged Maximum (% by weight). Sprouted or Insect Damaged.	2
of which, Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	1
Severely Damaged Max (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS Max (total count per 200 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8 falling below 4.33mm round hole screen	0.6
OTHER CONTAMINANTS Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Adzuki Bean seed material as outlined below.	1
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.1
Soil Maximum (% by weight)	n/a
Snails Maximum. Dead or alive. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects. Maximum Dead or alive per 200 gram sample.	15
Grasshoppers & Locusts Maximum Dead or alive per 200 gram sample	2
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-1.2 Adzuki Beans Minimum Export Standard Machine Dressed

(Grain Trade Australia) ADZUKI BEANS STANDARDS 2025/2026	
Commodity: Adzuki Beans	Season: 2025/2026
Grade: Minimum Export Standard Machine Dressed	Standard Reference No.: CSP 1.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The Adzuki Beans shall be sound, dry, fresh and have true varietal colour and characteristics for "Bloodwood" and "Erimo".	
Purity Minimum (% by weight). Whole Adzuki Beans, defective Adzuki Beans and Adzuki Beans seed coats.	99.5
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Adzuki Beans not of the specified variety and Adzuki Beans that are broken, chipped, damaged, diseased, frost damaged, insect damaged, loose seed coat, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed, and all Adzuki Bean seed material falling below the 4.76mm round hole screen.	2 Includes Severely Damaged
of which, Split/Broken Maximum (% by weight). Adzuki Beans that are not whole.	n/a
of which, Damaged Maximum (% by weight). Sprouted or Insect Damaged.	n/a
of which, Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	1
Severely Damaged Max (% by weight per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	Nil
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	Nil
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	Nil
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	Nil
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	Nil
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	Nil
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	Nil
Type 7a Max (seeds in total per 200 gram). Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	Nil
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	Nil
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	Nil
Type 8 Max (seeds in total per 200 gram). Bellvine	Nil
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8 falling below 4.33mm round hole screen	Nil
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Adzuki Bean seed material as outlined below.	0.5
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.1
Soil Maximum (% by weight)	n/a
Snails Maximum. Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects. Includes Grasshoppers and Locusts. Maximum Dead per 200 gram sample.	1
Of which Grasshoppers & Locusts Maximum Dead or alive per 200 gram sample	n/a
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-2.1.1 Broad Bean Minimum Receival Farmer Dressed

(Grain Trade Australia) BROAD BEANS STANDARDS 2025/2026	
Commodity: Broad Beans	Season: 2025/2026
Grade: Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP 2.1.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - Broad Beans shall be sound, dry, fresh and be colour typical for the variety of the season.	
Purity Minimum (% by weight). Whole Broad Beans, defective Broad Beans and Broad Beans seed coats.	97
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight in 600 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Broad Beans not of the specified variety and Broad Beans remaining above the 6mm slot screen that are broken, chipped, damaged, diseased, frost damaged, insect damaged, loose seed coat, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed, Kernel Damage, Poor Colour, Fungal Affected (e.g., Ascochyta) lesions and Screenings.	7
of which, Screenings Maximum (% by weight). All material passing through a 6 mm slotted screen is part of Screenings or Foreign Material	5
of which, Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class. Including evergreens (Max 2%), old season, dark beans, and nil tolerance for black beans.	3
of which, Fungal Affected (e.g., Ascochyta) lesions. Maximum (% by weight). Broad Beans affected with a spot greater than 4mm or more than one spot with combined size greater than 4mm. Affected is where the Fungal Affected (e.g., Ascochyta) lesion is visible on the seed coat.	3
of which, Insect Damaged Maximum (% by weight)	1.5
Severely Damaged Max (grains per 600 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS Max (total count per 600 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 600 gram. Except Type 1 in which the maximum applies on an individual seed basis per 600 gram.	
Type 1 Max* (Individual seeds per 600 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	12*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 600 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	3
Type 3b Max (seeds in total per 600 gram). Vetch (Commercial), Vetch (Tare)	6
Type 3c Max (seeds in total per 600 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	12
Type 4 Max (seeds in total per 600 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	30
Type 5 Max (seeds in total per 600 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	60
Type 6 Max (seeds in total per 600 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	15
Type 7a Max (seeds in total per 600 gram). Adzuki Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	30
Type 7b Max (seeds in total per 600 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Oats (Common), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	30
Type 7c Max (seeds in total per 600 gram). Safflower, Sunflower	3
Type 8 Max (seeds in total per 600 gram). Bellvine	300
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8 falling below 6mm slotted screen	0.6
OTHER CONTAMINANTS Max (% by weight in 600 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Broad Bean seed material as outlined below.	3
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum. Dead or alive. Whole or substantially whole (more than half) including bodies per 600g sample.	3
Field Insects. Maximum Dead or alive per 600g sample.	45
Grasshoppers & Locusts Maximum Dead or alive per 600g sample	6
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 600g sample.	6
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-2.1.2 Broad Beans Minimum Export Farmer Dressed

(Grain Trade Australia) BROAD BEANS STANDARDS 2025/2026	
Commodity: Broad Beans	Season: 2025/2026
Grade: Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP 2.1.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - Broad Beans shall be sound, dry, fresh and be colour typical for the variety of the season.	
Purity Minimum (% by weight). Whole Broad Beans, defective Broad Beans and Broad Beans seed coats.	97
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 600 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Broad Beans not of the specified variety and Broad Beans remaining above the 6mm slot screen that are broken, chipped, damaged, diseased, frost damaged, insect damaged, loose seed coat, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed, Kernel Damage, Poor Colour, Fungal Affected (e.g., Ascochyta) lesions and Screenings.	8 Includes Severely Damaged
of which, Screenings Maximum (% by weight). All material passing through a 6 mm slotted screen is part of Screenings or Foreign Material	5
of which, Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class. Including evergreens (Max 2%), old season, dark beans and black beans (1%).	3
of which, Fungal Affected (e.g., Ascochyta) lesions. Maximum (% by weight). Broad Beans affected with a spot greater than 4mm or more than one spot with combined size greater than 4mm. Affected is where the Fungal Affected (e.g., Ascochyta) lesion is visible on the seed coat.	3
of which, Insect Damaged Maximum (% by weight)	1.5
Severely Damaged Max (% by weight per 600 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 600 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 600 gram. Except Type 1 in which the maximum applies on an individual seed basis per 600 gram.	
Type 1 Max* (Individual seeds per 600 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	12*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 600 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	3
Type 3b Max (seeds in total per 600 gram). Vetch (Commercial), Vetch (Tare)	6
Type 3c Max (seeds in total per 600 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	12
Type 4 Max (seeds in total per 600 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	30
Type 5 Max (seeds in total per 600 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	60
Type 6 Max (seeds in total per 600 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	15
Type 7a Max (seeds in total per 600 gram). Adzuki Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	30
Type 7b Max (seeds in total per 600 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Oats (Common), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	30
Type 7c Max (seeds in total per 600 gram). Safflower, Sunflower	3
Type 8 Max (seeds in total per 600 gram). Bellvine	300
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8 falling below 6mm slotted screen	0.6
OTHER CONTAMINANTS - Max (% by weight in 600 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Broad Bean seed material as outlined below.	3
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum. Dead or alive. Whole or substantially whole (more than half) including bodies per 600 gram sample.	3
Field Insects. Maximum Dead or alive per 600 gram sample.	45
Grasshoppers & Locusts Maximum Dead or alive per 600 gram sample	6
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 600 gram sample.	6
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-2.1.3 Broad Beans Minimum Export Machine Dressed

(Grain Trade Australia) BROAD BEANS STANDARDS 2025/2026	
Commodity: Broad Beans	Season: 2025/2026
Grade: Minimum Export Standard Machine Dressed	Standard Reference No.: CSP 2.1.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - Broad Beans shall be sound, dry, fresh and be colour typical for the variety of the season.	
Purity Minimum (% by weight). Whole Broad Beans, defective Broad Beans and Broad Beans seed coats.	99.5
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 600 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Broad Beans not of the specified variety and Broad Beans remaining above the 6mm slot screen that are broken, chipped, damaged, diseased, frost damaged, insect damaged, loose seed coat, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed, Kernel Damage, Poor Colour, Fungal Affected (e.g., Ascochyta) lesions and Screenings.	7% Container 10% Bulk - includes Severely Damaged
of which, Screenings Maximum (% by weight). All material passing through a 6 mm slotted screen is part of Screenings or Foreign Material	1
of which, Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class. Including evergreens, old season, dark beans, and black beans.	3
of which, Fungal Affected (e.g., Ascochyta) lesions. Maximum (% by weight). Broad Beans affected with a spot greater than 4mm or more than one spot with combined size greater than 4mm. Affected is where the Fungal Affected (e.g., Ascochyta) lesion is visible on the seed coat.	3
of which, Insect Damaged Maximum (% by weight)	1.5
Severely Damaged Max (% by weight per 600g). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 600 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 600 gram. Except Type 1 in which the maximum applies on an individual seed basis per 600 gram.	
Type 1 Max* (Individual seeds per 600 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	12*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 600 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	3
Type 3b Max (seeds in total per 600 gram). Vetch (Commercial), Vetch (Tare)	6
Type 3c Max (seeds in total per 600 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	12
Type 4 Max (seeds in total per 600 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	30
Type 5 Max (seeds in total per 600 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	60
Type 6 Max (seeds in total per 600 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	15
Type 7a Max (seeds in total per 600 gram). Adzuki Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	30
Type 7b Max (seeds in total per 600 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Oats (Common), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	30
Type 7c Max (seeds in total per 600 gram). Safflower, Sunflower	3
Type 8 Max (seeds in total per 600 gram). Bellvine	300
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8 falling below 6mm slotted screen	0.6
OTHER CONTAMINANTS - Max (% by weight in 600 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Broad Bean seed material as outlined below.	0.5
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.1
Soil Maximum (% by weight)	
Snails Maximum. Dead or alive. Whole or substantially whole (more than half) including bodies per 600 gram sample.	Nil
Field Insects. Maximum Dead per 600 gram sample.	3 dead
Grasshoppers & Locusts Maximum Dead per 600 gram sample	
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 600 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch. The Broad Beans shall be designated into a category based on the final grade achieved by the majority (greater than 90%) of the Broad Beans once cleaned and graded e.g. 90% of Broad Beans must be retained above an 11mm round hole screen or 14mm round hole screen to be classified as 11mm or 14mm sized, respectively.	

CSP-2.2 Broad Beans No.1 Split Minimum Export Standard

(Grain Trade Australia) BROAD BEANS STANDARDS 2025/2026	
Commodity: Broad Beans	Season: 2025/2026
Grade: No.1 Split Minimum Export Standard	Standard Reference No.: CSP-2.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Broad Beans shall have a clean and bright appearance and shall be milled from hard and well-filled Broad Beans.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Split and whole Broad Beans seeds & caps but excludes detached Broad Beans seed coats.	99
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 600 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Broad Beans not of the specified variety and Broad Beans that are broken, chipped, damaged, diseased, frost damaged, insect damaged, sappy, shrivelled, sprouted, weather damaged, wrinkled, whole pods containing seed and immature Broad Beans, Caps, Broad Beans with the Seed Coat intact, Whole Broad Beans with a Missing Seed Coat and Poor Colour Broad Beans.	7 Includes Severely Damaged
of which, Poor Colour (Max % by weight) Broad Beans with green kernels from premature ripening and discoloured beans with cotyledons (seed) distinctly off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta) lesions.	3
of which, Broken and Kibbled (Max % by weight) Broad Bean seed material that passes through a 7.00 mm round hole screen	4
Of which, Caps (Max % by weight) Seed coats adhering to split or broken Broad Beans seed.	1
Severely Damaged Maximum (Max % by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 600 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 600 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 600 gram sample.	
Type 1 Max* (Individual seeds per 600 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	12*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 600 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	3
Type 3b Max (seeds in total per 600 gram). Vetch (Commercial), Vetch (Tare)	6
Type 3c Max (seeds in total per 600 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	12
Type 4 Max (seeds in total per 600 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	30
Type 5 Max (seeds in total per 600 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	60
Type 6 Max (seeds in total per 600 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	15
Type 7a Max (seeds in total per 600 gram). Adzuki Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	30
Type 7b Max (seeds in total per 600 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Oats (Common), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	30
Type 7c Max (seeds in total per 600 gram). Safflower, Sunflower	3
Type 8 Max (seeds in total per 600 gram). Bellvine	300
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8 falling below 6mm slotted screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 600 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Broad Bean seed material as outlined below. Includes detached seed coats.	1
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter.	0.1
Detached Seed Coats Maximum (% by weight)	0.1
Snails Maximum. Dead or alive. Whole or substantially whole (more than half) including bodies per 600 gram sample.	Nil
Field Insects. Maximum Dead or Alive per 600 gram sample.	Nil
Grasshoppers & Locusts Maximum Dead or Alive per 600 gram sample	Nil
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 600 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-3.1 Caloona / Poona Cowpeas Minimum Export Standard Machine Dressed

(Grain Trade Australia) CALOONA / POONA COWPEAS STANDARDS 2025/2026	
Commodity: Caloona / Poona Cowpeas	Season: 2025/2026
Grade: Minimum Export Standard Machine Dressed	Standard Reference No.: CSP 3.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The Cowpeas shall have a good bright appearance of the specified type i.e. buff coloured or red.	
Purity Minimum (% by weight). Whole Cowpeas, defective Cowpeas, Cowpeas other than specified type.	99
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Cowpeas not of the specified variety. Cowpeas that are broken, chipped, diseased, frost damaged, insect damaged, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Cowpeas, whether broken or unbroken and loose seed coat.	2
of which, Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta) lesions.	1
Severely Damaged Max (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8 falling below 4.33mm round hole screen	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Cowpea seed material as outlined below.	1
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.1
Soil Maximum (% by weight)	0.1
Snails Maximum. Dead or alive. Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects. Maximum Dead per 200 gram sample.	1 dead
Grasshoppers & Locusts Maximum Dead per 200 gram sample	1 dead
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
NOTE: Sowing seed shall have a minimum germination of 85% including hard seeds (I.S.T.A).	
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch. See Introduction.	

CSP-4.1.1 Chickpea - Desi No.1 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) CHICKPEA - DESI STANDARDS 2025/2026	
Commodity: Chickpea - Desi Grade: No.1 Minimum Receival Standard Farmer Dressed	Season: 2025/2026 Standard Reference No.: CSP- 4.1.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Desi type chickpeas should be sound, dry, fresh, and light to medium brown in colour (a slight greenish tinge of the seed coat is allowed). Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Includes whole Desi type chickpeas, defective Desi type chickpeas and seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Desi type chickpeas that are broken, chipped, fungal affected, frost damaged, diseased, green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained & weather damaged, wrinkled. Includes pods that contain Desi type chickpeas, whether broken or unbroken, loose seed coats and all Desi Chickpea seed material falling through the 3.97mm slotted screen.	6
of which, Poor Colour (% by weight) Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class, including green. Includes Fungal Affected (e.g., Ascochyta) lesions	2
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	1
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below 3.97mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Desi type chickpea seed material as outlined below.	3, includes Field Peas
Of which Field Peas Maximum (% by weight)	2
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non- vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds, and wood bugs. Whole only.	15
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-4.1.2 Chickpea - Desi No.1 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) CHICKPEA - DESI STANDARDS 2025/2026	
Commodity: Chickpea - Desi	Season: 2025/2026
Grade: No.1 Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP- 4.1.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Desi type chickpeas should be sound, dry, fresh, and light to medium brown in colour (a slight greenish tinge of the seed coat is allowed). Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Includes whole Desi type chickpeas, defective Desi type chickpeas and seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Desi type chickpeas that are broken, chipped, fungal affected, frost damaged, diseased, green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained & weather damaged, wrinkled. Includes pods that contain Desi type chickpeas, whether broken or unbroken, loose seed coats and all Desi Chickpea seed material falling through the 3.97mm slotted screen.	8% Cont. 10% Bulk Includes Severely Damaged
of which, Poor Colour (% by weight) Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class, including green. Includes Fungal Affected (e.g., Ascochyta) lesions	2
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	1
Severely Damaged Maximum (% by weight per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below 3.97mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Desi type chickpea seed material as outlined below.	3, includes Field Peas
Field Peas Maximum (% by weight)	2
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non- vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds, and wood bugs. Whole only.	15 dead
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-4.1.3 Chickpea - Desi No.1 Minimum Export Standard Machine Dressed

(Grain Trade Australia) CHICKPEA - DESI STANDARDS 2025/2026	
Commodity: Chickpea - Desi	Season: 2025/2026
Grade: No.1 Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-4.1.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Desi type chickpeas should be sound, dry, fresh, and light to medium brown in colour (a slight greenish tinge of the seed coat is allowed). Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Includes whole Desi type chickpeas, defective Desi type chickpeas and seed coats.	99
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Desi type chickpeas that are broken, chipped, fungal affected, frost damaged, diseased, green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained & weather damaged, wrinkled. Includes pods that contain Desi type chickpeas, whether broken or unbroken, loose seed coats and all Desi Chickpea seed material falling through the 3.97mm slotted screen.	3% Cont. 7% Bulk Includes Severely Damaged
of which, Poor Colour (% by weight) Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class, including green. Includes Fungal Affected (e.g., Ascochyta) lesions	2
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	1
Severely Damaged Maximum (% by weight per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below 3.97mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Desi type chickpea seed material as outlined below.	1
Field Peas Maximum (% by weight)	Included in Type 7a
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non-vegetable material.	0.1
Soil Maximum (% by weight)	0.1
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherghlen bugs, ladybirds and wood bugs. Whole only.	1 dead
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	1 dead
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-4.1.4 Chickpea - Desi No.2 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) CHICKPEA - DESI STANDARDS 2025/2026	
Commodity: Chickpea - Desi	Season: 2025/2026
Grade: No.2 Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP-4.1.4
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Desi type chickpeas should be sound, dry, fresh, and light to medium brown in colour (a slight greenish tinge of the seed coat is allowed). Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Includes whole Desi type chickpeas, defective Desi type chickpeas and seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Desi type chickpeas that are broken, chipped, fungal affected, frost damaged, diseased, green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained & weather damaged, wrinkled. Includes pods that contain Desi type chickpeas, whether broken or unbroken, loose seed coats and all Desi Chickpea seed material falling through the 3.97mm slotted screen.	10
of which, Poor Colour (% by weight) Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class, including green. Includes Fungal Affected (e.g., Ascochyta) lesions	5
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	1
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	10 ^
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below 3.97mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Desi type chickpea seed material as outlined below.	3, includes Field Peas
Of which Field Peas Maximum (% by weight)	2
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non- vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds, and wood bugs. Whole only.	15
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-4.1.5 Chickpea - Desi No.2 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) CHICKPEA - DESI STANDARDS 2025/2026	
Commodity: Chickpea - Desi	Season: 2025/2026
Grade: No.2 Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP-4.1.5
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Desi type chickpeas should be sound, dry, fresh, and light to medium brown in colour (a slight greenish tinge of the seed coat is allowed). Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Includes whole Desi type chickpeas, defective Desi type chickpeas and seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Desi type chickpeas that are broken, chipped, fungal affected, frost damaged, diseased, green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained & weather damaged, wrinkled. Includes pods that contain Desi type chickpeas, whether broken or unbroken, loose seed coats and all Desi Chickpea seed material falling through the 3.97mm slotted screen.	12% Cont. 15% Bulk Includes Severely Damaged
of which, Poor Colour (% by weight) Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class, including green. Includes Fungal Affected (e.g., Ascochyta) lesions	5
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	1
Severely Damaged Maximum (% by weight per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Bromegrass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below 3.97mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Desi type chickpea seed material as outlined below.	3, includes Field Peas
Of which Field Peas Maximum (% by weight)	2
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non- vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	15 dead
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-4.1.6 Chickpea - Desi No.3 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) CHICKPEA - DESI STANDARDS 2025/2026	
Commodity: Chickpea - Desi Grade: No.3 Minimum Receival Standard Farmer Dressed	Season: 2025/2026 Standard Reference No.: CSP- 4.1.6
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Desi type chickpeas should be sound, dry, fresh, and light to medium brown in colour (a slight greenish tinge of the seed coat is allowed). Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Includes whole Desi type chickpeas, defective Desi type chickpeas and seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Desi type chickpeas that are broken, chipped, fungal affected, frost damaged, diseased, green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained & weather damaged, wrinkled. Includes pods that contain Desi type chickpeas, whether broken or unbroken, loose seed coats and all Desi Chickpea seed material falling through the 3.97mm slotted screen.	15
of which, Poor Colour (% by weight) Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class, including green. Includes Fungal Affected (e.g., Ascochyta) lesions	n/a
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	n/a
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	15 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below 3.97mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Desi type chickpea seed material as outlined below.	3, includes Field Peas
Of which Field Peas Maximum (% by weight)	2
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non- vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds, and wood bugs. Whole only.	15
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-4.1.7 Chickpea - Desi No.3 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) CHICKPEA - DESI STANDARDS 2025/2026	
Commodity: Chickpea - Desi	Season: 2025/2026
Grade: No.3 Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP- 4.1.7
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Desi type chickpeas should be sound, dry, fresh, and light to medium brown in colour (a slight greenish tinge of the seed coat is allowed). Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Includes whole Desi type chickpeas, defective Desi type chickpeas and seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Desi type chickpeas that are broken, chipped, fungal affected, frost damaged, diseased, green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained & weather damaged, wrinkled. Includes pods that contain Desi type chickpeas, whether broken or unbroken, loose seed coats and all Desi Chickpea seed material falling through the 3.97mm slotted screen.	20% Cont. 25% Bulk Includes Severely Damaged
of which, Poor Colour (% by weight) Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class, including green. Includes Fungal Affected (e.g., Ascochyta) lesions	n/a
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	n/a
Severely Damaged Maximum (% by weight per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	2 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below 3.97mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Desi type chickpea seed material as outlined below.	3, includes Field Peas
Field Peas Maximum (% by weight)	2
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non- vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds, and wood bugs. Whole only.	15 dead
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-4.2 Chickpeas – Split Chana Dhal Minimum Export Standard

(Grain Trade Australia) CHICKPEA – SPLIT CHANA DHAL STANDARDS 2025/2026	
Commodity: Chickpea – Split Chana Dhal	Season: 2025/2026
Grade: Minimum Export Standard	Standard Reference No.: CSP-4.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Chana Dhal Split Chickpeas shall be milled from hard and well-filled chickpeas that are characteristic of the colour and variety.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Whole chickpeas with a missing seed coat, Split chickpeas, kibble and broken seed and greenish tinged and discoloured splits combined.	99
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Whole Chickpeas with a missing seed coat. Maximum (% by weight). Includes whole Chickpeas with a missing seed coat and Severely Damaged.	2
Of which Severely Damaged Maximum (% by weight per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
Poor Colour (Max % by weight) Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class, including green. Includes Fungal Affected (e.g., Ascochyta) lesions.	2
Fungal Affected (Max % by weight) Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel.	1
Broken and Kibbled (Max % by weight) Chickpea seed material which passes through a 3.57mm round hole screen.	4
Caps and Foreign Material (Max % by weight) Caps are the seed coats adhering to split or broken seed measured as a combined weight. Foreign material includes Unmillable Material, and all vegetable matter other than Desi type Chickpea seed material.	1
Detached Seed Coats (Max % by weight) Seed coats that are not attached to the kernels.	0.1
^ Not included in Total Defective	# Included in Whole chickpeas with a missing seed coat
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below 3.97mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Desi type chickpea seed material as outlined below. Includes Caps (Caps are the seed coats adhering to split or broken seed measured as a combined weight).	1
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non-vegetable material.	0.1
Soil Maximum (% by weight)	0.1
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	Nil
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	Nil
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-4.3.1 Chickpea Kabuli No.1 Large Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) CHICKPEA - KABULI STANDARDS 2025/2026	
Commodity: Chickpea Kabuli Grade: No.1 Large Minimum Receival Standard Farmer Dressed	Season: 2025/2026 Standard Reference No.: CSP-4.3.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Kabuli type chickpeas should be sound, dry, fresh, and cream to light brown in colour. Dark brown to Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Kabuli type Chickpeas, defective Kabuli type Chickpeas and seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in entire 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Kabuli Type Chickpeas not of the specified variety and Kabuli type Chickpeas that are broken, caked, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed and all Kabuli Chickpea seed material falling through the 6.00mm round hole screen. Includes poor colour.	3
of which, Poor Colour Maximum (% by weight). Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta) & Bleached grains.	2
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	1
Severely Damaged Maximum (grains per 400 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 400 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per entire 400 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram sample.	
Type 1 Max* (Individual seeds per 400 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Bromegrass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Oats (Common), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 6.00mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in entire 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Kabuli type Chickpeas seed material as outlined below.	3
Unmillable Material Maximum (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight).	0.3
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies, count in the 400 gram sample.	2
Field Insects Maximum Live or Dead count per 400 gram sample. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	30
Grasshoppers & Locusts Maximum Live or Dead. Count in total of the 400 gram sample. Whole or parts thereof.	4
Ryegrass Ergot Maximum (length in cm). Pieces laid end on end from the 400 gram sample.	4
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material. Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-4.3.2 Chickpea Kabuli No. 1 Large Minimum Export Standard Machine Dressed

(Grain Trade Australia) CHICKPEA - KABULI STANDARDS 2025/2026	
Commodity: Chickpea Kabul	Season: 2025/2026
Grade: No.1 Large Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-4.3.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS -The Kabuli type chickpeas should be sound, dry, fresh, and cream to light brown in colour. Dark brown to Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Kabuli type Chickpeas, defective Kabuli type Chickpeas and seed coats.	99.5
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight in entire 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Kabuli Type Chickpeas not of the specified variety and Kabuli type Chickpeas that are broken, caked, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed and all Kabuli Chickpea seed material falling through the 6.00mm round hole screen. Includes poor colour.	2 Includes Severely Damaged
of which, Poor Colour Maximum (% by weight). Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta) & Bleached grains.	2
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	1
Severely Damaged Maximum (% by weight per 400 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 400 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per entire 400 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per entire 400 gram sample.	
Type 1 Max* (Individual seeds per 400 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Oats (Common), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 6.00mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in entire 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Kabuli type Chickpeas seed material as outlined below.	0.5
Unmillable Material Maximum (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.1
Soil Maximum (% by weight).	Nil
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies, count in the 400 gram sample.	Nil
Field Insects Maximum Dead count per 400 gram sample. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	2 dead
Grasshoppers & Locusts Maximum Dead. Count in total of the 400 gram sample. Whole or parts thereof.	2 dead
Ryegrass Ergot Maximum (length in cm). Pieces laid end on end from the 400 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material. Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
The minimum and/or maximum size in millimetres applying to at least 92% of the Kabuli type chickpea of the lot agreed between the buyer and seller e.g. 8mm sized Kabuli type chickpeas, must have a retention of 92% above a 8mm round hole screen.	
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-4.3.3 Chickpea – Kabuli No.1 Small Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) CHICKPEA - KABULI STANDARDS 2025/2026	
Commodity: Chickpea Kabuli	Season: 2025/2026
Grade: No.1 Small Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP-4.3.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS -The Kabuli type chickpeas should be sound, dry, fresh, and cream to light brown in colour. Dark brown to Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Kabuli type Chickpeas, defective Kabuli type Chickpeas and seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS – Maximum (% by weight in entire 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Kabuli Type Chickpeas not of the specified variety and Kabuli type Chickpeas that are broken, caked, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed and all Kabuli Chickpea seed material falling through the 5.00mm round hole screen. Includes poor colour.	3
of which, Poor Colour Maximum (% by weight). Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta) & Bleached grains.	2
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	1
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS - Maximum (total count per entire 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per entire 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per entire 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 5.00mm screen	0.6
OTHER CONTAMINANTS Maximum (% by weight in entire 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Kabuli type Chickpeas seed material as outlined below.	3
Unmillable Material Maximum (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight).	0.3
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies, count in the 200 gram sample.	1
Field Insects Maximum Live or Dead count per 200 gram sample. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	15
Grasshoppers & Locusts Maximum Live or Dead. Count in total of the 200 gram sample. Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces laid end on end from the 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material. Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-4.3.4 Chickpea - Kabuli No.1 Small Minimum Export Standard Farmer Dressed

(Grain Trade Australia) CHICKPEA - KABULI STANDARDS 2025/2026	
Commodity: Chickpea - Kabuli	Season: 2025/2026
Grade: No.1 Small Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP-4.3.4
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Kabuli type chickpeas should be sound, dry, fresh, and cream to light brown in colour. Dark brown to Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Kabuli type Chickpeas, defective Kabuli type Chickpeas and seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in entire 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Kabuli Type Chickpeas not of the specified variety and Kabuli type Chickpeas that are broken, caked, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed and all Kabuli Chickpea seed material falling through the 5.00mm round hole screen. Includes poor colour.	8% cont. 10% bulk Includes Severely Damaged
of which, Poor Colour Maximum (% by weight). Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta) & Bleached grains.	2
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	1
Severely Damaged Maximum (% by weight per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per entire 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per entire 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per entire 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 5.00mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in entire 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Kabuli type Chickpeas seed material as outlined below.	3
Unmillable Material Maximum (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight).	0.3
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies, count in the 200 gram sample.	1
Field Insects Maximum Dead count per 200 gram sample. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	15
Grasshoppers & Locusts Maximum Dead. Count in total of the 200gram sample. Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces laid end on end from the 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material. Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-4.3.5 Chickpea – Kabuli No.1 Small Minimum Export Standard Machine Dressed

(Grain Trade Australia) CHICKPEA - KABULI STANDARDS 2025/2026	
Commodity: Chickpea Kabuli	Season: 2025/2026
Grade: No.1 Small Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-4.3.5
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS -The Kabuli type chickpeas should be sound, dry, fresh, and cream to light brown in colour. Dark brown to Black is excluded as the predominating class.	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Kabuli type Chickpeas, defective Kabuli type Chickpeas and seed coats.	99.5
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in entire 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Kabuli Type Chickpeas not of the specified variety and Kabuli type Chickpeas that are broken, caked, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed and all Kabuli Chickpea seed material falling through the 5.00mm round hole screen. Includes poor colour.	2 Includes Severely Damaged
of which, Poor Colour Maximum (% by weight). Kernel is distinctly blemished and / or off colour from the characteristic yellow colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta) & Bleached grains.	2
of which, Fungal Affected (e.g., Ascochyta) Maximum (% by weight). Fungal Affected (e.g., Ascochyta) means that a lesion is visible on the kernel. Classifiers are required to break the seed coat if they are not confident that the lesion has penetrated to the kernel.	1
Severely Damaged Maximum (% by weight per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per entire 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per entire 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per entire 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 5.00mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in entire 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Kabuli type Chickpeas seed material as outlined below.	0.5
Unmillable Material Maximum (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.1
Soil Maximum (% by weight).	
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies, count in the 200 gram sample.	Nil
Field Insects Maximum Dead count per 200 gram sample. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	1
Grasshoppers & Locusts Maximum Dead. Count in total of the 200gram sample. Whole or parts thereof.	
Ryegrass Ergot Maximum (length in cm). Pieces laid end on end from the 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material. Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
The minimum and/or maximum size in millimetres applying to at least 92% of the Kabuli type chickpea of the lot agreed between the buyer and seller e.g., 8mm sized Kabuli type chickpeas must have a retention of 92% above an 8mm round hole screen.	
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-5.1.1 Faba Beans – Canning Grade Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) FABA BEAN STANDARDS 2025/2026	
Commodity: Faba Beans	Season: 2025/2026
Grade: Canning Grade Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP-5.1.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Faba Beans should be sound, dry, and fresh and light to medium brown or pale green in colour	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Faba Beans, defective Faba Beans and Faba Bean seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Includes Faba Beans that are broken, chipped, diseased, frost damaged, fully green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained, weather damaged, wrinkled. Includes pods that contain Faba Beans, whether broken or unbroken, loose seed coats and all Faba Bean seed material falling through the 3.75mm slotted screen	2
of which, Poor Colour Maximum (% by weight). Seed Coat is distinctly blemished and / or off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta).	1
Retention Minimum (% by weight). 90% of Faba Beans must not pass through the screen to be classified as 8mm.	90%
Severely Damaged Maximum (grains per 400 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
# Included in Total Defective	^ Not included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 400 gram; unless otherwise stated). Tolerances apply to whole seeds or their equivalent in pieces and refer to the max total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1- 8 falling below 3.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes unmillable material and all vegetable matter other than Faba Bean seed material. Foreign Material includes stalks and plant material that may be connected to the plant and all the below parameters.	3
Unmillable Material Maximum (% by weight). Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Live or Dead. Whole or substantially whole (more than half) including bodies count per 400 gram sample.	2
Field Insects. Live or Dead (count per 400 grams). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Siton Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	30
Grasshoppers & Locusts. Live or dead (count per 400 gram). Whole or parts thereof.	4
Ryegrass Ergot (length in cm). Pieces laid end to end per 400 gram sample.	4
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-5.1.2 Faba Beans – Canning Grade Minimum Export Standard Machine Dressed

(Grain Trade Australia) FABA BEAN STANDARDS 2025/2026	
Commodity: Faba Beans	Season: 2025/2026
Grade: Canning Grade Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-5.1.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Faba Beans should be sound, dry and fresh and light to medium brown or pale green in colour	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Faba Beans, defective Faba Beans and Faba Bean seed coats.	99.5
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Includes Faba Beans that are broken, chipped, diseased, frost damaged, fully green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained, weather damaged, wrinkled. Includes pods that contain Faba Beans, whether broken or unbroken, loose seed coats and all Faba Bean seed material falling through the 3.75mm slotted screen	3
of which, Poor Colour Maximum (% by weight). Seed Coat is distinctly blemished and / or off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta).	1
of which Severely Damaged Maximum (% by weight per 400 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1% #
Retention Minimum (% by weight). 90% of Faba Beans must not pass through the screen to be classified as 8mm.	90%
# Included in Total Defective	^ Not included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 400 gram; unless otherwise stated). Tolerances apply to whole seeds or their equivalent in pieces and refer to the max total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1- 8 falling below 3.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes unmillable material and all vegetable matter other than Faba Bean seed material. Foreign Material includes stalks and plant material that may be connected to the plant and all the below parameters.	0.5
Unmillable Material Maximum (% by weight). Includes soil, stones, sclerotes and other non-vegetable material.	0.1
Soil Maximum (% by weight)	
Snails Live or Dead. Whole or substantially whole (more than half) including bodies count per 400 gram sample.	Nil
Field Insects. Dead (count per 400 grams). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	2
Grasshoppers & Locusts. Dead (count per 400 gram). Whole or parts thereof.	
Ryegrass Ergot (length in cm). Pieces laid end to end per 400 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
The Faba beans shall be designated into a category based on the final grade achieved by the majority (greater than 90%) once cleaned and graded e.g., 90% of faba beans must be retained above an 8mm round hole screen to be classified as 8mm.	
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-5.2.1 Faba Beans No.1 Minimum Receival Farmer Dressed

(Grain Trade Australia) FABA BEAN STANDARDS 2025/2026	
Commodity: Faba Beans	Season: 2025/2026
Grade: No.1 Minimum Receival Farmer Dressed	Standard Reference No.: CSP-5.2.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Faba Beans should be sound, dry, and fresh and light to medium brown or pale green in colour	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Faba Beans, defective Faba Beans and Faba Bean seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Includes Faba Beans that are broken, chipped, diseased, frost damaged, fully green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained, weather damaged, wrinkled. Includes pods that contain Faba Beans, whether broken or unbroken, loose seed coats and all Faba Bean seed material falling through the 3.75mm slotted screen	6
of which, Poor Colour Maximum (% by weight). Seed Coat is distinctly blemished and / or off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta).	4
of which, Burnt, Caked, Heat Damaged, Sprouted Maximum (% by weight)	N/A
Severely Damaged Maximum (grains per 400 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
Mould (field or storage) Maximum (grains per 400 gram)	N/A
# Included in Total Defective	^ Not included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 400 gram; unless otherwise stated). Tolerances apply to whole seeds or their equivalent in pieces and refer to the max total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1- 8 falling below 3.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Faba Bean seed material. Foreign Material includes stalks and plant material that may be connected to the plant and all the below parameters.	3
Unmillable Material Maximum (% by weight). Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Live or Dead. Whole or substantially whole (more than half) including bodies count per 400 gram sample.	2
Field Insects. Live or Dead (count per 400 grams). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Siton Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	30
Grasshoppers & Locusts. Live or dead (count per 400 gram). Whole or parts thereof.	4
Ryegrass Ergot (length in cm). Pieces laid end to end per 400 gram sample.	4
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-5.2.2 Faba Beans - No.1 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) FABA BEAN STANDARDS 2025/2026	
Commodity: Faba Beans	Season: 2025/2026
Grade: No.1 Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP-5.2.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Faba Beans should be sound, dry and fresh and light to medium brown or pale green in colour	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Faba Beans, defective Faba Beans and Faba Bean seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Includes Faba Beans that are broken, chipped, diseased, frost damaged, fully green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained, weather damaged, wrinkled. Includes pods that contain Faba Beans, whether broken or unbroken, loose seed coats and all Faba Bean seed material falling through the 3.75mm slotted screen	8% cont. 10% bulk
of which, Poor Colour Maximum (% by weight). Seed Coat is distinctly blemished and / or off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta).	4
of which, Burnt, Caked, Heat Damaged, Sprouted Maximum (% by weight)	N/A
Severely Damaged Maximum (% by weight per 400 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
Mould (field or storage) Maximum (grains per 400 gram)	N/A
# Included in Total Defective	^ Not included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 400 gram; unless otherwise stated). Tolerances apply to whole seeds or their equivalent in pieces and refer to the max total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1- 8 falling below 3.75mm screen	0.6
OTHER CONTAMINANTS Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Faba Bean seed material. Foreign Material includes stalks and plant material that may be connected to the plant and all the below parameters.	3
Unmillable Material Maximum (% by weight). Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails. Dead. Whole or substantially whole (more than half) including bodies count per 400 gram sample.	2
Field Insects. Dead (count per 400 grams). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	30
Grasshoppers & Locusts. Dead (count per 400 gram). Whole or parts thereof.	4
Ryegrass Ergot (length in cm). Pieces laid end to end per 400 gram sample.	4
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch	

CSP-5.2.3 Faba Beans No.1 Minimum Export Standard Machine Dressed

(Grain Trade Australia) FABA BEAN STANDARDS 2025/2026	
Commodity: Faba Beans	Season: 2025/2026
Grade: No.1 Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-5.2.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Faba Beans should be sound, dry and fresh and light to medium brown or pale green in colour	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Faba Beans, defective Faba Beans and Faba Bean seed coats.	99
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Includes Faba Beans that are broken, chipped, diseased, frost damaged, fully green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained, weather damaged, wrinkled. Includes pods that contain Faba Beans, whether broken or unbroken, loose seed coats and all Faba Bean seed material falling through the 3.75mm slotted screen	6% cont. 10% bulk
of which, Poor Colour Maximum (% by weight). Seed Coat is distinctly blemished and / or off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta).	4
of which, Burnt, Caked, Heat Damaged, Sprouted Maximum (% by weight)	N/A
Severely Damaged Maximum (% by weight per 400 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
Mould (field or storage) Maximum (grains per 400 gram)	N/A
# Included in Total Defective	^ Not included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 400 gram; unless otherwise stated). Tolerances apply to whole seeds or their equivalent in pieces and refer to the max total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1- 8 falling below 3.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Faba Bean seed material. Foreign Material includes stalks and plant material that may be connected to the plant and all the below parameters.	1
Unmillable Material Maximum (% by weight). Includes soil, stones, sclerotes and other non-vegetable material.	0.1
Soil Maximum (% by weight)	0.1
Snails Dead. Whole or substantially whole (more than half) including bodies count per 400 gram sample.	Nil
Field Insects. Dead (count per 400 grams). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	2
Grasshoppers & Locusts. Dead (count per 400 gram). Whole or parts thereof.	2
Ryegrass Ergot (length in cm). Pieces laid end to end per 400 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch	

CSP-5.3.1 Faba Beans - No.2 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) FABA BEAN STANDARDS 2025/2026	
Commodity: Faba Beans Grade: No.2 Minimum Receival Standard Farmer Dressed	Season: 2025/2026 Standard Reference No.: CSP-5.3.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS The Faba Beans should be sound, dry, and fresh and light to medium brown or pale green in colour	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Faba Beans, defective Faba Beans and Faba Bean seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Includes Faba Beans that are broken, chipped, diseased, frost damaged, fully green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained, weather damaged, wrinkled. Includes pods that contain Faba Beans, whether broken or unbroken, loose seed coats and all Faba Bean seed material falling through the 3.75mm slotted screen	10
of which, Poor Colour Maximum (% by weight). Seed Coat is distinctly blemished and / or off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta).	7
of which, Burnt, Caked, Heat Damaged, Sprouted Maximum (% by weight)	N/A
Severely Damaged Maximum (grains per 400 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
Mould (field or storage) Maximum (grains per 400 gram)	N/A
# Included in Total Defective	^ Not included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 400 gram; unless otherwise stated). Tolerances apply to whole seeds or their equivalent in pieces and refer to the max total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1- 8 falling below 3.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Faba Bean seed material. Foreign Material includes stalks and plant material that may be connected to the plant and all the below parameters.	3
Unmillable Material Maximum (% by weight). Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Live or Dead. Whole or substantially whole (more than half) including bodies count per 400 gram sample.	2
Field Insects. Live or Dead (count per 400 grams). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	30
Grasshoppers & Locusts. Live or dead (count per 400 gram). Whole or parts thereof.	4
Ryegrass Ergot (length in cm). Pieces laid end to end per 400 gram sample.	4
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-5.3.2 Faba Beans - No.2 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) FABA BEAN STANDARDS 2025/2026	
Commodity: Faba Beans	Season: 2025/2026
Grade: No.2 Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP-5.3.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS The Faba Beans should be sound, dry, and fresh and light to medium brown or pale green in colour	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Faba Beans, defective Faba Beans and Faba Bean seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS Maximum (% by weight, 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Includes Faba Beans that are broken, chipped, diseased, frost damaged, fully green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained, weather damaged, wrinkled. Includes pods that contain Faba Beans, whether broken or unbroken, loose seed coats and all Faba Bean seed material falling through the 3.75mm slotted screen	12% cont. 14% bulk
of which, Poor Colour Maximum (% by weight). Seed Coat is distinctly blemished and / or off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta).	7
of which, Burnt, Caked, Heat Damaged, Sprouted Maximum (% by weight)	N/A
Severely Damaged Maximum (% by weight per 400 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
Mould (field or storage) Maximum (grains per 400 gram)	N/A
# Included in Total Defective	^ Not included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 400 gram; unless otherwise stated). Tolerances apply to whole seeds or their equivalent in pieces and refer to the max total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1- 8 falling below 3.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Faba Bean seed material. Foreign Material includes stalks and plant material that may be connected to the plant and all the below parameters.	3
Unmillable Material Maximum (% by weight). Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Dead. Whole or substantially whole (more than half) including bodies count per 400 gram sample.	2
Field Insects. Dead (count per 400 grams). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	30
Grasshoppers & Locusts. Dead (count per 400 gram). Whole or parts thereof.	4
Ryegrass Ergot (length in cm). Pieces laid end to end per 400 gram sample.	4
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch	

CSP-5.4.1 Faba Beans - No.3 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) FABA BEAN STANDARDS 2025/2026	
Commodity: Faba Beans	Season: 2025/2026
Grade: No.3 Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP-5.4.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS The Faba Beans should be sound, dry, and fresh and light to medium brown or pale green in colour	
Moisture (%) Maximum	14
Purity Minimum (% by weight). Whole Faba Beans, defective Faba Beans and Faba Bean seed coats.	97
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Includes Faba Beans that are broken, chipped, diseased, frost damaged, fully green, hail damaged, insect damaged, poor colour, sappy, shrivelled, split, sprouted, stained, weather damaged, wrinkled. Includes pods that contain Faba Beans, whether broken or unbroken, loose seed coats and all Faba Bean seed material falling through the 3.75mm slotted screen	20
of which, Poor Colour Maximum (% by weight). Seed Coat is distinctly blemished and / or off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta).	N/A
of which, Burnt, Caked, Heat Damaged, Sprouted Maximum (% by weight)	7
Severely Damaged Maximum (grains per 400 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	N/A
Mould (field or storage) Maximum (grains per 400 gram)	1 ^
# Included in Total Defective	^ Not included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 400 gram; unless otherwise stated). Tolerances apply to whole seeds or their equivalent in pieces and refer to the max total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1- 8 falling below 3.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Faba Bean seed material. Foreign Material includes stalks and plant material that may be connected to the plant and all the below parameters.	3
Unmillable Material Maximum (% by weight). Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Live or Dead. Whole or substantially whole (more than half) including bodies count per 400 gram sample.	2
Field Insects. Live or Dead (count per 400 grams). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	30
Grasshoppers & Locusts. Live or dead (count per 400 gram). Whole or parts thereof.	4
Ryegrass Ergot (length in cm). Pieces laid end to end per 400 gram sample.	4
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	1
Objectionable Material (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-5.5 Faba Beans – No.1 Split Minimum Export Standard Machine Dressed

(Grain Trade Australia) FABA BEANS STANDARDS 2025/2026	
Commodity: Faba Beans	Season: 2025/2026
Grade: No.1 Split Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-5.5
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - Split Faba Beans having a clean bright appearance. The Faba Beans shall be milled from hard and well filled whole Faba Beans.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Includes Whole Faba Beans, split Faba Beans, defective Faba Beans and Faba Bean seed coats.	99.5
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Tolerances apply to Poor Colour, Caps and Faba Beans with the Seed Coat intact, Faba Beans with a Missing Seed Coat and Severely Damaged.	3
Of which Poor Colour (Max % by weight) Kernel which is distinctly off colour from the characteristic colour of the predominating class. Includes Fungal Affected (e.g., Ascochyta) and Green Faba Beans.	2#
Of which Caps and Faba Beans with Seed Coat Intact (Max % by weight) Caps are seed coats adhering to split or broken seed.	2#
Of which Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1#
Broken and Kibbled (Max % by weight) Faba Bean seed material that passes through a 6mm round hole screen.	4^
Detached Seed Coats (Max % by weight) Seed coats that are not attached to the kernels.	0.1 ^
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 400 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 400 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram sample.	
Type 1 Max* (Individual seeds per 400 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1- 8 falling below 3.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Faba Bean seed material as outlined below.	0.5
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non-vegetable material. Maximum of 1 Stone per kg sample.	0.1
Soil Maximum (% by weight)	
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 400 gram sample.	Nil
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	Nil
Grasshoppers & Locusts Maximum Live or Dead per 400 gram sample. Whole or parts thereof.	
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 400 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-6.1 Fenugreek - Whole – No.1 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) FENUGREEK STANDARDS 2025/2026	
Commodity: Fenugreek	Season: 2025/2026
Grade: Whole No.1 Minimum Receival Farmer Dressed	Standard Reference No.: CSP 6.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS – The Fenugreek shall be hard and well filled.	
Purity Minimum (% by weight). Whole Fenugreek, defective Fenugreek and Fenugreek Seed Coats.	97
Moisture (%) Maximum	12
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Fenugreek not of the specified variety and Fenugreek that is broken, chipped, damaged, diseased, frost damaged, insect damaged, loose seed coat, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed, Kernel Damage, Poor Colour, Fungal Affected (e.g., Ascochyta) lesions and all Fenugreek seed material falling through the 1.4mm slotted screen.	3
of which, Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class. Includes poor colour arising from disease such as bacterial blight, frost damage and water staining.	1
of which, Varietal Restriction. Maximum (% by weight). Fenugreek not of the specified variety.	1
Severely Damaged Max (grains per 200g). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS Max (total count per 200 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 1.4mm slotted screen	0.6
OTHER CONTAMINANTS Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Fenugreek seed material as outlined below.	3
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum. Dead or alive. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects. Maximum Dead or alive per 200 gram sample.	15
Grasshoppers & Locusts Maximum Dead or alive per 200 gram sample	2
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-6.2 Fenugreek - Whole – No.1 Minimum Export Standard Machine Dressed

(Grain Trade Australia) FENUGREEK STANDARDS 2025/2026	
Commodity: Fenugreek	Season: 2025/2026
Grade: Whole No.1 Minimum Export Standard Machine Dressed	Standard Reference No.: CSP 6.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The Fenugreek shall be hard and well filled.	
Purity Minimum (% by weight). Whole Fenugreek, defective Fenugreek and Fenugreek Seed Coats.	99
Moisture (%) Maximum	12
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Fenugreek not of the specified variety and Fenugreek that is broken, chipped, damaged, diseased, frost damaged, insect damaged, loose seed coat, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes whole pods containing seed, Kernel Damage, Poor Colour, Fungal Affected (e.g., Ascochyta) lesions and all Fenugreek seed material falling through the 1.4mm slotted screen.	2% Includes Severely Damaged
of which, Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class. Includes poor colour arising from disease such as bacterial blight, frost damage and water staining.	1
of which, Varietal Restriction. Maximum (% by weight). Fenugreek not of the specified variety.	1
Severely Damaged Max (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 1.4mm slotted screen	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Fenugreek seed material as outlined below.	1
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.1
Soil Maximum (% by weight)	0.1
Snails Maximum. Dead or alive. Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects. Maximum Dead or alive per 200 gram sample.	Nil
Grasshoppers & Locusts Maximum Dead or alive per 200 gram sample	Nil
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-7.1.1 Lentils – Whole Green No.1 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) LENTILS – WHOLE GREEN STANDARDS 2025/2026	
Commodity: Green Lentils	Season: 2025/2026
Grade: Whole Green No.1 Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP-7.1.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Green Lentils should be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Moisture (%) Maximum	14
Purity (% by weight) Minimum. Includes whole Green Lentils, defective Green Lentils and Green Lentil Seed Coats	97
DEFECTIVE GRAINS - Max (% by weight, 200 gram sample; unless otherwise stated)	
Total Defective (% by weight) Lentils not of the specified variety. Lentil kernels that are black, broken, chipped, diseased, frost damaged, insect damaged, orange tip, sappy, shrivelled, split, sprouted, weather damaged, wrinkled, poor colour seed coat and poor colour kernel. Includes pods that contain Green Lentils, whether broken or unbroken, loose seed coat, and all Green Lentil seed material falling through the slotted 2.0mm screen (Boomer 2.2mm).	4.0
of which Varietal Restriction (% by weight) Maximum. Green Lentils not of the specified variety	1.0
of which Poor Colour Seed Coat (% by weight) Maximum. Discoloured Green Lentils have excessive discolouration of the seed coat. Includes any disease, frost, and water staining. Includes Fungal Affected (e.g., Ascochyta)	1.0
of which Poor Colour Kernel (% by weight) Maximum. Discoloured Green Lentils have excessive discolouration of the kernel. Includes any disease, frost and water staining, and green, brown, black, yellow, bleached, and chalky white kernels. Includes Fungal Affected (e.g., Ascochyta)	1.0
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Cowpea, Faba Beans, Lentils (Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram) Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Ryegrass on Stalk, Turnip Weed Pods. Any other seed not specified in TYPE (1) to TYPE(7a), TYPE 7(c), TYPE 8, CEREAL SEEDS or SFS	10
Type 7c Max (seeds in total per 200 gram) Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram) Bellvine	100
Cereal Seeds (seeds in total per 200 gram) Barley, Corn (Maize), Durum, Oats (Common), Rice, Rye (Cereal), Sorghum, Triticale, Wheat	2
Small Foreign Seeds Max (% by weight) Foreign seeds not specified in TYPES 1-8 or CEREAL SEEDS falling below the specified screen.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum Includes unmillable material and all vegetable matter other than Green Lentil seed material as outlined below.	3
of which Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
of which Soil (% by weight) Maximum	0.3
Snails Live or Dead (count per 200 gram) Maximum. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Live or Dead (count per 200 gram) Maximum. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	15
Grasshoppers, Locusts Live or Dead (count per 200 gram) Maximum. Whole or parts thereof.	2
Ryegrass Ergot Max (length in cm per 200 gram)	2.0
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Max (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell or taste.	Nil

CSP-7.1.2 Lentils – Whole Green No.1 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) LENTILS – WHOLE GREEN STANDARDS 2025/2026	
Commodity: Green Lentils	Season: 2025/2026
Grade: Whole Green No.1 Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP-7.1.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Green Lentils should be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Moisture (%) Maximum	14
Purity (% by weight) Minimum. Includes whole Green Lentils, defective Green Lentils and Green Lentil Seed Coats	97
DEFECTIVE GRAINS - Max (% by weight, 200 gram sample; unless otherwise stated)	
Total Defective (% by weight) Green Lentils not of the specified variety. Green Lentil kernels that are black, broken, chipped, diseased, frost damaged, insect damaged, orange tip, sappy, shrivelled, split, sprouted, weather damaged, wrinkled, poor colour seed coat and poor colour kernel. Includes pods that contain Green Lentils, whether broken or unbroken, loose seed coat, and all Green Lentil seed material falling through the slotted 2.0mm screen (Boomer 2.2mm).	6% cont. 10% bulk
of which Varietal Restriction (% by weight) Maximum. Green Lentils not of the specified variety	1.0
of which Poor Colour Seed Coat (% by weight) Maximum. Discoloured Green Lentils have excessive discolouration of the seed coat. Includes any disease, frost, and water staining. Includes Fungal Affected (e.g., Ascochyta)	1.0
of which Poor Colour Kernel (% by weight) Maximum. Discoloured Green Lentils have excessive discolouration of the kernel. Includes any disease, frost and water staining, and green, brown, black, yellow, bleached, and chalky white kernels. Includes Fungal Affected (e.g., Ascochyta)	1.0
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Cowpea, Faba Beans, Lentils (Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram) Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Ryegrass on Stalk, Turnip Weed Pods. Any other seed not specified in TYPE (1) to TYPE(7a), TYPE 7(c), TYPE 8, CEREAL SEEDS or SFS	10
Type 7c Max (seeds in total per 200 gram) Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram) Bellvine	100
Cereal Seeds (seeds in total per 200 gram) Barley, Corn (Maize), Durum, Oats (Common), Rice, Rye (Cereal), Sorghum, Triticale, Wheat	2
Small Foreign Seeds Max (% by weight) Foreign seeds not specified in TYPES 1-8 or CEREAL SEEDS falling below the specified screen.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes unmillable material and all vegetable matter other than Green Lentil seed material as outlined below.	3
of which Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
of which Soil (% by weight) Maximum	0.3
Snails Live or Dead (count per 200 gram) Maximum. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Live or Dead (count per 200 gram) Maximum. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds, and wood bugs. Whole only.	15
Grasshoppers, Locusts Live or Dead (count per 200 gram) Maximum. Whole or parts thereof.	2
Ryegrass Ergot Max (length in cm per 200 gram)	2.0
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Max (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-7.1.3 Lentils – Whole Green No.1 Minimum Export Standard Machine Dressed

(Grain Trade Australia) LENTILS – WHOLE GREEN STANDARDS 2025/2026	
Commodity: Green Lentils	Season: 2025/2026
Grade: Whole Green No.1 Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-7.1.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Green Lentils should be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Moisture (%) Maximum	14
Purity (% by weight) Minimum. Includes whole Green Lentils, defective Green Lentils and Green Lentil Seed Coats	99
DEFECTIVE GRAINS - Max (% by weight, 200 gram sample; unless otherwise stated)	
Total Defective (% by weight). Green Lentils not of the specified variety. Green Lentil kernels that are black, broken, chipped, diseased, frost damaged, insect damaged, orange tip, sappy, shrivelled, split, sprouted, weather damaged, wrinkled, poor colour seed coat and poor colour kernel. Includes pods that contain Green Lentils, whether broken or unbroken, loose seed coat, and all Green Lentil seed material falling through the slotted 2.0mm screen (Boomer 2.2mm).	3% cont. 5% bulk
of which Varietal Restriction (% by weight) Maximum. Green Lentils not of the specified variety	1.0
of which Poor Colour Seed Coat (% by weight) Maximum. Discoloured Green Lentils have excessive discolouration of the seed coat. Includes any disease, frost, and water staining. Includes Fungal Affected (e.g., Ascochyta)	1.0
of which Poor Colour Kernel (% by weight) Maximum. Discoloured Green Lentils have excessive discolouration of the kernel. Includes any disease, frost and water staining, and green, brown, black, yellow, bleached, and chalky white kernels. Includes Fungal Affected (e.g., Ascochyta)	1.0
Severely Damaged Maximum (& by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Cowpea, Faba Beans, Lentils (Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram) Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Ryegrass on Stalk, Turnip Weed Pods. Any other seed not specified in TYPE (1) to TYPE (7a), TYPE 7(c), TYPE 8, CEREAL SEEDS or SFS	10
Type 7c Max (seeds in total per 200 gram) Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram) Bellvine	100
Cereal Seeds (seeds in total per 200 gram) Barley, Corn (Maize), Durum, Oats (Common), Rice, Rye (Cereal), Sorghum, Triticale, Wheat	2
Small Foreign Seeds Max (% by weight) Foreign seeds not specified in TYPES 1-8 or CEREAL SEEDS falling below the specified screen.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable material and all vegetable matter other than Green Lentil seed material as outlined below	1
of which Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.1
of which Soil (% by weight) Maximum	
Snails Live or Dead (count per 200 gram) Maximum. Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects Live or Dead (count per 200 gram) Maximum Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	1 dead
Grasshoppers, Locusts Live or Dead (count per 200 gram) Maximum. Whole or parts thereof.	
Ryegrass Ergot Max (length in cm per 200 gram)	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Max (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-7.2.1 Lentil - Whole Red No.1 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) LENTILS – WHOLE RED STANDARDS 2025/2026	
Commodity: Red Lentils	Season: 2025/2026
Grade: Whole Red No.1 Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP-7.2.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Red Lentils should be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Moisture (%) Maximum	14
Purity (% by weight) Minimum. Includes whole Red Lentils, defective Red Lentils and Red Lentil Seed Coats	97
DEFECTIVE GRAINS - Max (% by weight, 200 gram sample; unless otherwise stated)	
Total Defective (% by weight) Red Lentils not of the specified variety. Red Lentil kernels that are black, broken, chipped, diseased, frost damaged, insect damaged, orange tip, sappy, shrivelled, split, sprouted, weather damaged, wrinkled, poor colour seed coat and poor colour kernel. Includes pods that contain Red Lentils, whether broken or unbroken, loose seed coat, 3% blondes (Aldinga only) and all Red Lentil seed material falling through the slotted 2.0mm screen (Aldinga 2.2mm).	4.0
of which Varietal Restriction (% by weight) Maximum. Red Lentils not of the specified variety	1.0
of which Poor Colour Seed Coat (% by weight) Maximum. Discoloured Red Lentils have excessive discolouration of the seed coat as per the GTA Visual Recognition Standards Guide. Includes any disease, frost, and water staining. Includes Fungal Affected (e.g., Ascochyta)	1.0
of which Poor Colour Kernel (% by weight) Maximum. Discoloured Red Lentils have excessive discolouration of the kernel as per the GTA Visual Recognition Standards Guide. Includes any disease, frost and water staining, and green, brown, black, yellow, bleached, and chalky white kernels. Includes Fungal Affected (e.g., Ascochyta)	1.0
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
Contrasting Colour (% by weight) Maximum. Contrasting colour of the seed coat in the same variety. Not classified as defective.	1.0 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Cowpea, Faba Beans, Lentils (Green), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram) Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Ryegrass on Stalk, Turnip Weed Pods. Any other seed not specified in TYPE (1) to TYPE(7a), TYPE 7(c), TYPE 8, CEREAL SEEDS or SFS	10
Type 7c Max (seeds in total per 200 gram) Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram) Bellvine	100
Cereal Seeds (seeds in total per 200 gram) Barley, Corn (Maize), Durum, Oats (Common), Rice, Rye (Cereal), Sorghum, Triticale, Wheat	2
Small Foreign Seeds Max (% by weight) Foreign seeds not specified in TYPES 1-8 or CEREAL SEEDS falling below the specified screen.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable material and all vegetable matter other than Red Lentil seed material as outlined below	3
of which Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
of which Soil (% by weight) Maximum	0.3
Snails Live or Dead (count per 200 gram) Maximum. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Live or Dead (count per 200 gram) Maximum. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	15
Grasshoppers, Locusts Live or Dead (count per 200 gram) Maximum. Whole or parts thereof.	2
Ryegrass Ergot Max (length in cm per 200 gram)	2.0
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Max (entire load) Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell or taste.	Nil

CSP-7.2.2 Lentils – Whole Red No.1 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) LENTILS – WHOLE RED STANDARDS 2025/2026	
Commodity: Red Lentils	Season: 2025/2026
Grade: Whole Red No.1 Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP-7.2.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Red Lentils should be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Moisture (%) Maximum	14
Purity (% by weight) Minimum. Includes whole Red Lentils, defective Red Lentils and Red Lentil Seed Coats	97
DEFECTIVE GRAINS - Max (% by weight, 200 gram sample; unless otherwise stated)	
Total Defective (% by weight). Red Lentils not of the specified variety. Red Lentil kernels that are black, broken, chipped, diseased, frost damaged, insect damaged, orange tip, sappy, shrivelled, split, sprouted, weather damaged, wrinkled, poor colour seed coat and poor colour kernel. Includes pods that contain Red Lentils, whether broken or unbroken, loose seed coat, 3% blondes (Aldinga only) and all Red Lentil seed material falling through the slotted 2.0mm screen (Aldinga 2.2mm).	6% cont. 10% bulk
of which Varietal Restriction (% by weight) Maximum. Red Lentils not of the specified variety	1.0
of which Poor Colour Seed Coat (% by weight) Maximum. Discoloured Red Lentils have excessive discolouration of the seed coat as per the GTA Visual Recognition Standards Guide. Includes any disease, frost, and water staining. Includes Fungal Affected (e.g., Ascochyta)	1.0
of which Poor Colour Kernel (% by weight) Maximum. Discoloured Red Lentils have excessive discolouration of the kernel as per the GTA Visual Recognition Standards Guide. Includes any disease, frost and water staining, and green, brown, black, yellow, bleached, and chalky white kernels. Includes Fungal Affected (e.g., Ascochyta)	1.0
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
Contrasting Colour (% by weight) Maximum. Contrasting colour of the seed coat in the same variety. Not classified as defective.	1.0 ^
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Cowpea, Faba Beans, Lentils (Green), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram) Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Ryegrass on Stalk, Turnip Weed Pods. Any other seed not specified in TYPE (1) to TYPE (7a), TYPE 7(c), TYPE 8, CEREAL SEEDS or SFS	10
Type 7c Max (seeds in total per 200 gram) Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram) Bellvine	100
Cereal Seeds (seeds in total per 200 gram) Barley, Corn (Maize), Durum, Oats (Common), Rice, Rye (Cereal), Sorghum, Triticale, Wheat	2
Small Foreign Seeds Max (% by weight) Foreign seeds not specified in TYPES 1-8 or CEREAL SEEDS falling below the specified screen.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable material and all vegetable matter other than Red Lentil seed material as outlined below	3
of which Unmillable Material (% by weight) Maximum Includes soil, stones, sclerotes and other non-vegetable material.	0.5
of which Soil (% by weight) Maximum	0.3
Snails Dead (count per 200 gram) Maximum Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Dead (count per 200 gram) Maximum Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds, and wood bugs. Whole only.	15
Grasshoppers, Locusts Live or Dead (count per 200 gram) Maximum. Whole or parts thereof.	2
Ryegrass Ergot Max (length in cm per 200 gram)	2.0
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Max (entire load) Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract.	
Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-7.2.3 Lentils – Whole Red No.1 Minimum Export Standard Machine Dressed

(Grain Trade Australia) LENTILS – WHOLE RED STANDARDS 2025/2026	
Commodity: Red Lentils	Season: 2025/2026
Grade: Whole Red No.1 Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-7.2.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Red Lentils should be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Moisture (%) Maximum	14
Purity (% by weight) Minimum. Includes whole Red Lentils, defective Red Lentils and Red Lentil Seed Coats	99
DEFECTIVE GRAINS - Max (% by weight, 200 gram sample; unless otherwise stated)	
Total Defective (% by weight) Red Lentils not of the specified variety. Red Lentil kernels that are black, broken, chipped, diseased, frost damaged, insect damaged, orange tip, sappy, shrivelled, split, sprouted, weather damaged, wrinkled, poor colour seed coat and poor colour kernel. Includes pods that contain Red Lentils, whether broken or unbroken, loose seed coat, 3% blondes (Aldinga only) and all Red Lentil seed material falling through the slotted 2.0mm screen (Aldinga 2.2mm).	3% cont. 5% bulk
of which Varietal Restriction (% by weight) Maximum. Red Lentils not of the specified variety	1.0
of which Poor Colour Seed Coat (% by weight) Maximum. Discoloured Red Lentils have excessive discoloration of the seed coat as per the GTA Visual Recognition Standards Guide. Includes any disease, frost, and water staining. Includes Fungal Affected (e.g., Ascochyta)	1.0
of which Poor Colour Kernel (% by weight) Maximum. Discoloured Red Lentils have excessive discoloration of the kernel as per the GTA Visual Recognition Standards Guide. Includes any disease, frost and water staining, and green, brown, black, yellow, bleached, and chalky white kernels. Includes Fungal Affected (e.g., Ascochyta)	1.0
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
Contrasting Colour (% by weight) Maximum. Contrasting colour of the seed coat in the same variety. Not classified as defective.	1.0 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Cowpea, Faba Beans, Lentils (Green), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram) Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Ryegrass on Stalk, Turnip Weed Pods. Any other seed not specified in TYPE (1) to TYPE (7a), TYPE 7(c), TYPE 8, CEREAL SEEDS or SFS	10
Type 7c Max (seeds in total per 200 gram) Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram) Bellvine	100
Cereal Seeds (seeds in total per 200 gram) Barley, Corn (Maize), Durum, Oats (Common), Rice, Rye (Cereal), Sorghum, Triticale, Wheat	2
Small Foreign Seeds Max (% by weight) Foreign seeds not specified in TYPES 1-8 or CEREAL SEEDS falling below the specified screen.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum Includes Unmillable material and all vegetable matter other than Red Lentil seed material as outlined below	1
of which Unmillable Material (% by weight) Maximum Includes soil, stones, sclerotes and other non-vegetable material.	0.1
of which Soil (% by weight) Maximum	0.1
Snails Live or Dead (count per 200 gram) Maximum Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects Dead (count per 200 gram) Maximum Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	1
Grasshoppers, Locusts Dead (count per 200 gram) Maximum. Whole or parts thereof.	1
Ryegrass Ergot Max (length in cm per 200 gram)	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Max (entire load) Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell or taste.	Nil

When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.

CSP-7.3.1 Lentils - Whole Red No.2 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) LENTILS – WHOLE RED STANDARDS 2025/2026	
Commodity: Red Lentils	Season: 2025/2026
Grade: Whole Red No.2 Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP-7.3.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Red Lentils should be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Moisture (%) Maximum	14
Purity (% by weight) Minimum. Includes whole Red Lentils, defective Red Lentils and Red Lentil Seed Coats	97
DEFECTIVE GRAINS - Max (% by weight, 200 gram sample; unless otherwise stated)	
Total Defective (% by weight) Red Lentils not of the specified variety. Red Lentil kernels that are black, broken, chipped, diseased, frost damaged, insect damaged, orange tip, sappy, shrivelled, split, sprouted, weather damaged, wrinkled, poor colour seed coat and poor colour kernel. Includes pods that contain Red Lentils, whether broken or unbroken, loose seed coat, 3% blondes (Aldinga only) and all Red Lentil seed material falling through the slotted 2.0mm screen (Aldinga 2.2mm).	8.0
of which Varietal Restriction (% by weight) Maximum. Red Lentils not of the specified variety	1.0
of which Poor Colour Seed Coat (% by weight) Maximum. Discoloured Red Lentils have excessive discolouration of the seed coat as per the GTA Visual Recognition Standards Guide. Includes any disease, frost, and water staining. Includes Fungal Affected (e.g., Ascochyta)	3.0
of which Poor Colour Kernel (% by weight) Maximum. Discoloured Red Lentils have excessive discolouration of the kernel as per the GTA Visual Recognition Standards Guide. Includes any disease, frost and water staining, and green, brown, black, yellow, bleached and chalky white kernels. Includes Fungal Affected (e.g., Ascochyta)	1.0
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
Contrasting Colour (% by weight) Maximum. Contrasting colour of the seed coat in the same variety. Not classified as defective.	1.0 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS Max (total count per 200 gram; unless otherwise stated) Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Cowpea, Faba Beans, Lentils (Green), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram) Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Ryegrass on Stalk, Turnip Weed Pods. Any other seed not specified in TYPE (1) to TYPE(7a), TYPE 7(c), TYPE 8, CEREAL SEEDS or SFS	10
Type 7c Max (seeds in total per 200 gram) Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram) Bellvine	100
Cereal Seeds (seeds in total per 200 gram) Barley, Corn (Maize), Durum, Oats (Common), Rice, Rye (Cereal), Sorghum, Triticale, Wheat	2
Small Foreign Seeds Max (% by weight) Foreign seeds not specified in TYPES 1-8 or CEREAL SEEDS falling below the specified screen.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum Includes Unmillable material and all vegetable matter other than Red Lentil seed material as outlined below	3
of which Unmillable Material (% by weight) Maximum Includes soil, stones, sclerotes and other non-vegetable material.	0.5
of which Soil (% by weight) Maximum	0.3
Snails Live or Dead (count per 200 gram) Maximum Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Live or Dead (count per 200 gram) Maximum Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	15
Grasshoppers, Locusts Live or Dead (count per 200 gram) Maximum. Whole or parts thereof.	2
Ryegrass Ergot Max (length in cm per 200 gram)	2.0
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Max (entire load) Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell or taste.	Nil

CSP-7.3.2 Lentils Whole Red No.2 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) LENTILS – WHOLE RED STANDARDS 2025/2026	
Commodity: Red Lentils	Season: 2025/2026
Grade: Whole Red No.2 Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP-7.3.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Red Lentils should be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Moisture (%) Maximum	14
Purity (% by weight) Minimum. Includes whole Red Lentils, defective Red Lentils and Red Lentil Seed Coats	97
DEFECTIVE GRAINS - Max (% by weight, 200 gram sample; unless otherwise stated)	
Total Defective (% by weight) Red Lentils not of the specified variety. Red Lentil kernels that are black, broken, chipped, diseased, frost damaged, insect damaged, orange tip, sappy, shrivelled, split, sprouted, weather damaged, wrinkled, poor colour seed coat and poor colour kernel. Includes pods that contain Red Lentils, whether broken or unbroken, loose seed coat, 3% blondes (Aldinga only) and all Red Lentil seed material falling through the slotted 2.0mm screen (Aldinga 2.2mm).	9% cont. 14% bulk
of which Varietal Restriction (% by weight) Maximum. Red Lentils not of the specified variety	1.0
of which Poor Colour Seed Coat (% by weight) Maximum. Discoloured Red Lentils have excessive discolouration of the seed coat as per the GTA Visual Recognition Standards Guide. Includes any disease, frost, and water staining. Includes Fungal Affected (e.g., Ascochyta)	3.0
of which Poor Colour Kernel (% by weight) Maximum. Discoloured Red Lentils have excessive discolouration of the kernel as per the GTA Visual Recognition Standards Guide. Includes any disease, frost and water staining, and green, brown, black, yellow, bleached and chalky white kernels. Includes Fungal Affected (e.g., Ascochyta)	1.0
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
Contrasting Colour (% by weight) Maximum. Contrasting colour of the seed coat in the same variety. Not classified as defective.	1.0 ^
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Cowpea, Faba Beans, Lentils (Green), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram) Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Ryegrass on Stalk, Turnip Weed Pods. Any other seed not specified in TYPE (1) to TYPE (7a), TYPE 7(c), TYPE 8, CEREAL SEEDS or SFS	10
Type 7c Max (seeds in total per 200 gram) Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram) Bellvine	100
Cereal Seeds (seeds in total per 200 gram) Barley, Corn (Maize), Durum, Oats (Common), Rice, Rye (Cereal), Sorghum, Triticale, Wheat	2
Small Foreign Seeds Max (% by weight) Foreign seeds not specified in TYPES 1-8 or CEREAL SEEDS falling below the specified screen.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable material and all vegetable matter other than Red Lentil seed material as outlined below	3
of which Unmillable Material (% by weight) Maximum Includes soil, stones, sclerotes and other non-vegetable material.	0.5
of which Soil (% by weight) Maximum	0.3
Snails Dead (count per 200 gram) Maximum Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Dead (count per 200 gram) Maximum Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	15
Grasshoppers, Locusts Live or Dead (count per 200 gram) Maximum. Whole or parts thereof.	2
Ryegrass Ergot Max (length in cm per 200 gram)	2.0
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Max (entire load) Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-7.3.3 Lentil Whole Red No.2 Minimum Export Standard Machine Dressed

(Grain Trade Australia) LENTILS – WHOLE RED STANDARDS 2025/2026	
Commodity: Red Lentils	Season: 2025/2026
Grade: Whole Red No.2 Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-7.3.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - The Red Lentils should be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Moisture (%) Maximum	14
Purity (% by weight) Minimum. Includes whole Red Lentils, defective Red Lentils and Red Lentil Seed Coats	99
DEFECTIVE GRAINS - Max (% by weight, 200 gram sample; unless otherwise stated)	
Total Defective (% by weight). Red Lentils not of the specified variety. Red Lentil kernels that are black, broken, chipped, diseased, frost damaged, insect damaged, orange tip, sappy, shrivelled, split, sprouted, weather damaged, wrinkled, poor colour seed coat and poor colour kernel. Includes pods that contain Red Lentils, whether broken or unbroken, loose seed coat, 3% blondes (Aldinga only) and all Red Lentil seed material falling through the slotted 2.0mm screen (Aldinga 2.2mm).	7% cont. 9% bulk
of which Varietal Restriction (% by weight) Maximum. Red Lentils not of the specified variety	1.0
of which Poor Colour Seed Coat (% by weight) Maximum. Discoloured Red Lentils have excessive discolouration of the seed coat as per the GTA Visual Recognition Standards Guide. Includes any disease, frost, and water staining. Includes Fungal Affected (e.g., Ascochyta)	3.0
of which Poor Colour Kernel (% by weight) Maximum. Discoloured Red Lentils have excessive discolouration of the kernel as per the GTA Visual Recognition Standards Guide. Includes any disease, frost and water staining, and green, brown, black, yellow, bleached, and chalky white kernels. Includes Fungal Affected (e.g., Ascochyta)	1.0
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
Contrasting Colour (% by weight) Maximum. Contrasting colour of the seed coat in the same variety. Not classified as defective.	1.0 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS Max (total count per 200 gram; unless otherwise stated) Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Cowpea, Faba Beans, Lentils (Green), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram) Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Ryegrass on Stalk, Turnip Weed Pods. Any other seed not specified in TYPE (1) to TYPE(7a), TYPE 7(c), TYPE 8, CEREAL SEEDS or SFS	10
Type 7c Max (seeds in total per 200 gram) Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram) Bellvine	100
Cereal Seeds (seeds in total per 200 gram) Barley, Corn (Maize), Durum, Oats (Common), Rice, Rye (Cereal), Sorghum, Triticale, Wheat	2
Small Foreign Seeds Max (% by weight) Foreign seeds not specified in TYPES 1-8 or CEREAL SEEDS falling below the specified screen.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum Includes Unmillable material and all vegetable matter other than Red Lentil seed material as outlined below	1
of which Unmillable Material (% by weight) Maximum Includes soil, stones, sclerotes and other non-vegetable material.	0.1
of which Soil (% by weight) Maximum	
Snails Live or Dead (count per 200 gram) Maximum Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects Dead (count per 200 gram) Maximum Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	1
Grasshoppers, Locusts Dead (count per 200 gram) Maximum. Whole or parts thereof.	
Ryegrass Ergot Max (length in cm per 200 gram)	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Max (entire load) Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-7.4.1 Lentils Split Red No.1 Minimum Export Standard

(Grain Trade Australia) LENTILS – SPLIT RED STANDARDS 2025/2026	
Commodity: Red Lentils	Season: 2025/2026
Grade: Split Red No.1 Minimum Export Standard	Standard Reference No.: CSP-7.4.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - Split Red Lentils having a clean and bright appearance. The Lentils shall be milled from hard and well-filled whole Red Lentil seed.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Split Red Lentils, whole Red Lentils, Broken & Kibbled and Caps.	99.75
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Poor Colour (Max % by weight) Red Lentil kernel that is distinctly off colour from the characteristic colour of the predominating class. Includes kernels that are brown, black, green, yellow or bleached as per the GTA Visual Recognition Standards Guide.	0.25
Chalky White Disease (Max % by weight) Kernels that have a distinct chalky white lesion as per the GTA Visual Recognition Standards Guide.	0.25
Caps (Max % by weight) Caps are seed coats adhering to split or broken seed.	0.25
Whole Red Lentils with a Missing Seed Coat (Max % by weight) Whole Red Lentils with a missing seed coat	5
Broken and Kibbled (Max % by weight) Red Lentil seed material that passes through a 3mm round hole screen (Northfield 2.78mm).	5
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the specified screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Red Lentil seed material including detached seed coats.	0.25
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non-vegetable material.	
Soil Maximum (% by weight)	
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	Nil
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	Nil
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-7.4.2 Lentils Split Red No.2 Minimum Export Standard

(Grain Trade Australia) LENTILS – SPLIT RED STANDARDS 2025/2026	
Commodity: Red Lentils	Season: 2025/2026
Grade: Split Red No.2 Minimum Export Standard	Standard Reference No.: CSP-7.4.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - Split Red Lentils having a clean and bright appearance. The Lentils shall be milled from hard and well-filled whole Red Lentil seed	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Split Red Lentils, whole Red Lentils, Broken & Kibbled and Caps.	99.5
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Poor Colour (Max % by weight) Red Lentil Kernel that is distinctly off colour from the characteristic colour of the predominating class. Includes kernels that are brown, black, green, yellow or bleached as per the GTA Visual Recognition Standards Guide.	0.25
Chalky White Disease (Max % by weight) Kernels that have a distinct chalky white lesion as per the GTA Visual Recognition Standards Guide.	0.25
Caps (Max % by weight) Caps are seed coats adhering to split or broken seed.	0.5
Whole Red Lentils with a Missing Seed Coat (Max % by weight) Whole Red Lentils with a missing seed coat	5
Broken and Kibbled (Max % by weight) Red Lentil seed material that passes through a 3mm round hole screen (Northfield 2.78mm).	5
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the specified screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Red Lentil seed material including detached seed coats.	0.5
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non-vegetable material.	
Soil Maximum (% by weight)	
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	Nil
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	Nil
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-7.4.3 Lentils Split Red No.3 Minimum Export Standard

(Grain Trade Australia) LENTILS – SPLIT RED STANDARDS 2025/2026	
Commodity: Red Lentils	Season: 2025/2026
Grade: Split Red No.3 Minimum Export Standard	Standard Reference No.: CSP-7.4.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - Split Red Lentils having a clean and bright appearance. The Lentils shall be milled from hard and well-filled whole Red Lentil seed.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Split Red Lentils, whole Red Lentils, Broken & Kibbled and Caps.	99
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Poor Colour (Max % by weight) Red Lentil Kernel that is distinctly off colour from the characteristic colour of the predominating class. Includes kernels that are brown, black, green, yellow or bleached as per the GTA Visual Recognition Standards Guide.	0.25
Poor Colour Yellow (Max % by weight). Yellow kernel that is distinctly off colour from the characteristic colour of the predominating class as per the GTA Visual Recognition Standards Guide.	2
Chalky White Disease (Max % by weight) Kernels that have a distinct chalky white lesion as per the GTA Visual Recognition Standards Guide.	0.25
Caps (Max % by weight) Caps are seed coats adhering to split or broken seed.	2
Whole Red Lentils with a Missing Seed Coat (Max % by weight) Whole Red Lentils with a missing seed coat	5
Broken and Kibbled (Max % by weight) Red Lentil seed material that passes through a 3mm round hole screen (Northfield 2.78mm).	5
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the specified screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Red Lentil seed material including detached seed coats.	1
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non-vegetable material.	
Soil Maximum (% by weight)	
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	Nil
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	Nil
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-8.1.1 Lupins Angustifolius Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) LUPINS – ANGUSTIFOLIUS STANDARDS 2025/2026	
Commodity: Angustifolius Lupins	Season: 2025/2026
Grade: Angustifolius Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP – 8.1.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The Angustifolius Lupins shall be of the current season and be dry and mature.	
Purity Minimum (% by weight). Whole Angustifolius Lupins, defective Angustifolius Lupins and Angustifolius Lupin seed coats.	97
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Angustifolius Lupins not of the specified type. Angustifolius Lupins that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Angustifolius Lupins, whether broken or unbroken, loose seed coats, poor colour, bitter dark seeded lupins, Phomopsis affected and all Angustifolius Lupin seed material falling through the 3.75mm slotted screen.	7
of which, Poor Colour (count per 200 gram) Maximum. Yellow reddish / tan coloured Angustifolius Lupins as per the GTA Visual Recognition Standards Guide.	36
of which, Bitter Dark Seeded Lupins Maximum (count per 200 gram)	2
of which, Phomopsis Affected Lupins Maximum (count per 200 gram)	17
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling through the 3.75mm slotted screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in entire 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Angustifolius Lupin seed material as outlined below. Includes 2% tolerance for wild radish.	3
Unmillable Material (% by weight) Maximum. Includes soil, stones, metal and other non-vegetable material.	0.5
Soil Maximum (% by weight).	0.3
Snails. Dead or alive. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Maximum count per 200 gram. Dead or alive. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	15
Grasshoppers & Locusts Maximum dead or alive count per 200 gram. Whole or parts thereof.	2
Ryegrass Ergot (length in cm) Pieces aligned end on end per 200 gram.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-8.1.3 Lupins Angustifolius Minimum Export Standard Farmer Dressed

(Grain Trade Australia) LUPINS – ANGUSTIFOLIUS STANDARDS 2025/2026	
Commodity: Angustifolius Lupins Grade: Angustifolius Minimum Export Standard Farmer Dressed	Season: 2025/2026 Standard Reference No.: CSP-8.1.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The Angustifolius Lupins shall be of the current season and be dry and mature.	
Purity Minimum (% by weight). Whole Angustifolius Lupins, defective Angustifolius Lupins and Angustifolius Lupin seed coats.	97
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Angustifolius Lupins not of the specified type. Angustifolius Lupins that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Angustifolius Lupins, whether broken or unbroken, loose seed coats, poor colour, bitter dark seeded lupins Phomopsis affected and all Angustifolius Lupin seed material falling through the 3.75mm slotted screen.	9% Cont. 11% Bulk
of which, Poor Colour (count per 200 gram) Maximum. Yellow reddish / tan coloured Angustifolius Lupins as per the GTA Visual Recognition Standards Guide.	36
of which, Bitter Dark Seeded Lupins Maximum (count per 200 gram)	2
of which, Phomopsis Affected Lupins Maximum (count per 200 gram)	17
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 3.75mm slotted screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in entire 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Angustifolius Lupin seed material as outlined below. Includes 2% tolerance for wild radish.	3
Unmillable Material (% by weight) Maximum. Includes soil, stones, metal and other non-vegetable material.	0.5
Soil Maximum (% by weight).	0.3
Snails. Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	1
Field Insects Maximum count per 200 gram. Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only) Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	15
Grasshoppers & Locusts Maximum dead or alive count per 200 gram. Whole or parts thereof.	2
Ryegrass Ergot (length in cm) Pieces aligned end on end per 200 gram.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-8.1.4 Lupins Angustifolius Minimum Export Standard Machine Dressed

(Grain Trade Australia) LUPINS – ANGUSTIFOLIUS STANDARDS 2025/2026	
Commodity: Angustifolius Lupins	Season: 2025/2026
Grade: Angustifolius Minimum Export Standard Machine Dressed	Standard Reference No.: CSP – 8.1.4
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The Angustifolius Lupins shall be of the current season and be dry and mature.	
Purity Minimum (% by weight). Whole Angustifolius Lupins, defective Angustifolius Lupins and Angustifolius Lupin seed coats.	99.5
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Angustifolius Lupins not of the specified type. Angustifolius Lupins that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Angustifolius Lupins, whether broken or unbroken, loose seed coats, poor colour, bitter dark seeded lupins, Phomopsis affected and all Angustifolius Lupin seed material falling through the 3.75mm slotted screen.	4
of which, Poor Colour (count per 200 gram) Maximum. Yellow reddish / tan coloured Angustifolius Lupins as per the GTA Visual Recognition Standards Guide.	36
of which, Bitter Dark Seeded Lupins Maximum (count per 200 gram)	2
of which, Phomopsis Affected Lupins Maximum (count per 200 gram)	17
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 3.75mm slotted screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in entire 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Angustifolius Lupin seed material as outlined below. Includes 2% tolerance for wild radish.	0.5
Unmillable Material (% by weight) Maximum. Includes soil, stones, metal and other non-vegetable material.	0.1
Soil Maximum (% by weight).	0.1
Snails. Dead or alive. Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects Maximum count per 200 gram. Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	1
Grasshoppers & Locusts Maximum dead count per 200 gram. Whole or parts thereof.	1
Ryegrass Ergot (length in cm) Pieces aligned end on end per 200 gram.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-8.2.1 Lupins – Albus No.1 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) LUPINS – ALBUS STANDARDS 2025/2026	
Commodity: Albus Lupins Grade: Albus No.1 Minimum Receival Standard Farmer Dressed	Season: 2025/2026 Standard Reference No.: CSP – 8.2.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The Albus Lupins shall be of the current season and be dry and mature.	
Purity (% by weight) Minimum. Whole Albus Lupins, Defective Albus Lupins and Albus Lupin seed coats.	97
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Albus Lupins not of the specified variety. Albus Lupins that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Albus Lupins, whether broken or unbroken, loose seed coat, poor colour, bitter dark seeded lupins, Phomopsis affected and all Albus Lupin seed material falling through the 6.75m round hole screen	5
of which, Poor Colour (% by weight) Maximum. Albus Lupins whose seed coat or kernels are distinctly off colour from the characteristic colour of the predominating class.	1
of which, Bitter Dark Seeded Lupins Maximum (count per 400 gram)	4
of which, Phomopsis Affected Seeds Maximum (count per 400 gram)	34
Severely Damaged Maximum (grains per 400 grams). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 [^]
[^] Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 400 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 6.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Albus Lupin seed material as outlined below. Includes 2% tolerance for wild radish.	3
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails. Dead or Alive Maximum Whole or substantially whole (more than half) including bodies count per 400 gram sample.	2
Field Insects Maximum Dead or Alive count per 400 gram. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	30
Grasshoppers & Locusts Maximum Dead or Alive (count per 400 gram). Whole or parts thereof.	4
Ryegrass Ergot Maximum (length in cm per 400 gram). Pieces aligned end on end.	4
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire sample). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-8.2.2 Lupins – Albus No.1 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) LUPINS – ALBUS STANDARDS 2025/2026	
Commodity: Albus Lupins	Season: 2025/2026
Grade: Albus No. 1 Minimum Export Standard Farmer Dressed	Standards Reference No.: CSP- 8.2.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The Albus Lupins shall be of the current season and be dry and mature.	
Purity (% by weight) Minimum. Whole Albus Lupins, Defective Albus Lupins and Albus Lupin seed coats.	97
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Albus Lupins not of the specified variety. Albus Lupins that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Albus Lupins, whether broken or unbroken, loose seed coat, poor colour, bitter dark seeded lupins, Phomopsis affected and all Albus Lupin seed material falling through the 6.75mm round hole screen	5
of which, Poor Colour (% by weight) Maximum. Albus Lupins whose seed coat or kernels are distinctly off colour from the characteristic colour of the predominating class.	1
of which, Bitter Dark Seeded Lupins Maximum (count per 400 gram)	4
of which, Phomopsis Affected Seeds Maximum (count per 400 gram)	34
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 400 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 6.75mm screen	0.6
OTHER CONTAMINANTS Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Albus Lupin seed material as outlined below. Includes 2% tolerance for wild radish.	3
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails. Dead. Maximum Whole or substantially whole (more than half) including bodies count per 400 gram sample.	2
Field Insects Maximum Dead. Count per 400 gram. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	30
Grasshoppers & Locusts Maximum Dead (count per 400 gram). Whole or parts thereof.	4
Ryegrass Ergot Maximum (length in cm per 400 gram). Pieces aligned end on end.	4
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire sample). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-8.2.3 Lupins – Albus No.1 Minimum Export Standard Machine Dressed

(Grain Trade Australia) LUPINS – ALBUS STANDARDS 2025/2026	
Commodity: Albus Lupins Grade: Albus No. 1 Minimum Export Standard Machine Dressed	Season: 2025/2026 Standard Reference No.: CSP – 8.2.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The Albus Lupins shall be of the current season and be dry and mature.	
Purity (% by weight) Minimum. Whole Albus Lupins, Defective Albus Lupins and Albus Lupin seed coats.	99.5
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Albus Lupins not of the specified variety. Albus Lupins that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Albus Lupins, whether broken or unbroken, loose seed coat, poor colour, bitter dark seeded lupins, Phomopsis affected and all Albus Lupin seed material falling through the 6.75mm round hole screen	2
of which, Poor Colour (% by weight) Maximum. Albus Lupins whose seed coat or kernels are distinctly off colour from the characteristic colour of the predominating class.	1
of which, Bitter Dark Seeded Lupins Maximum (count per 400 gram)	4
of which, Phomopsis Affected Seeds Maximum (count per 400 gram)	34
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS -Maximum (total count per 400 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Catrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 6.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Albus Lupin seed material as outlined below. Includes 2% tolerance for wild radish.	0.5
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.1
Soil Maximum (% by weight)	0.1
Snails. Dead or Alive Maximum Whole or substantially whole (more than half) including bodies count per 400 gram sample.	Nil
Field Insects Maximum Dead (count per 400 gram). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	2
Grasshoppers & Locusts Maximum Dead (count per 400 gram). Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm per 400 gram). Pieces aligned end on end.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Maximum (entire sample). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-8.3.1 Lupins – Albus No.2 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) LUPINS – ALBUS STANDARDS 2025/2026	
Commodity: Albus Lupins Grade: Albus No. 2 Minimum Receival Standard Farmer Dressed	Season: 2025/2026 Standards Reference No.: CSP- 8.3.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The Albus Lupins shall be of the current season and be dry and mature.	
Purity (% by weight) Minimum. Whole Albus Lupins, Defective Albus Lupins and Albus Lupin seed coats.	97
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Albus Lupins not of the specified variety. Albus Lupins that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Albus Lupins, whether broken or unbroken, loose seed coat, poor colour, bitter dark seeded lupins, Phomopsis affected and all Albus Lupin seed material falling through the 6.75mm round hole screen	7
of which, Poor Colour (% by weight) Maximum. Albus Lupins whose seed coat or kernels are distinctly off colour from the characteristic colour of the predominating class.	2
of which, Bitter Dark Seeded Lupins Maximum (count per 400 gram)	4
of which, Phomopsis Affected Seeds Maximum (count per 400 gram)	34
Severely Damaged Maximum (grains per 400 grams). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 [^]
[^] Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 400 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 400 gram. Except Type 1 in which the maximum applies on an individual seed basis per 400 gram.	
Type 1 Max* (Individual seeds per 400 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	8*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 400 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	2
Type 3b Max (seeds in total per 400 gram). Vetch (Commercial), Vetch (Tare)	4
Type 3c Max (seeds in total per 400 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	8
Type 4 Max (seeds in total per 400 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20
Type 5 Max (seeds in total per 400 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	40
Type 6 Max (seeds in total per 400 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	10
Type 7a Max (seeds in total per 400 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	20
Type 7b Max (seeds in total per 400 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	20
Type 7c Max (seeds in total per 400 gram). Safflower, Sunflower	2
Type 8 Max (seeds in total per 400 gram). Bellvine	200
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below the 6.75mm screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 400 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Albus Lupin seed material as outlined below. Includes 2% tolerance for wild radish.	3
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails. Dead or Alive Maximum Whole or substantially whole (more than half) including bodies count per 400 gram sample.	2
Field Insects Maximum Dead or Alive count per 400 gram. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	30
Grasshoppers & Locusts Maximum Dead or Alive (count per 400 gram). Whole or parts thereof.	4
Ryegrass Ergot Maximum (length in cm per 400 gram). Pieces aligned end on end.	4
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire sample). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-10.1.1 Peas – Field No.1 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) PEAS – FIELD STANDARDS 2025/2026	
Commodity: Peas - Field Grade: No.1 Minimum Receival Standard Farmer Dressed	Season: 2025/2026 Standard Reference No. CSP- 10.1.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The peas shall be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Purity (% by weight) Minimum. Whole Field Peas, Defective Field Peas and Field Pea seed coats.	97
Moisture (%) Maximum	14
+ NOTE: For the No.1 grade, not less than 70% by weight of the entire sample shall be prime peas, that is, field peas of a size that will not pass through a 6.35mm round hole screen.	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Field Peas not of the specified variety. Field Peas that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Field Peas, whether broken or unbroken, loose seed coat and all Field Pea seed material falling through the 3.75m slotted screen. Includes Poor Colour.	3 +
of which, Poor Colour (% by weight) Maximum. Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	1
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective # Is included in Total Defective + Note: Screen only used for assessment of No.1 grade	
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8 (including falling below the 3.75mm screen where required to be used).	0.6
Clover Burr Maximum (count per 200 gram). Included in Foreign Material	1
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable Material and all vegetable matter other than Field Pea seed material as outlined below	3
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Dead or Alive (count per 200 gram). Whole or substantially whole (more than half) including bodies	1
Field Insects Maximum Dead or Alive (count per 200 gram). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole.	15
Grasshoppers & Locusts Maximum Dead or Alive (count per 200 gram). Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces aligned end on end	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-10.1.2 Peas – Field No.1 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) PEAS – FIELD STANDARDS 2025/2026	
Commodity: Peas - Field	Season: 2025/2026
Grade: No.1 Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP- 10.1.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The peas shall be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Purity (% by weight) Minimum. Whole Field Peas, Defective Field Peas and Field Pea seed coats.	97
Moisture (%) Maximum	14
+ NOTE: For the No.1 grade, not less than 70% by weight of the entire sample shall be prime peas, that is, field peas of a size that will not pass through a 6.35mm round hole screen.	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Field Peas not of the specified variety. Field Peas that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Field Peas, whether broken or unbroken, loose seed coat and all Field Pea seed material falling through the 3.75m slotted screen. Includes Poor Colour.	5% cont. + 7% bulk +
of which, Poor Colour (% by weight) Maximum. Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	1
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Is included in Total Defective
+ Note: Screen only used for assessment of No.1 grade	
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8 (including falling below the 3.75mm screen where required to be used).	0.6
Clover Burr Maximum (count per 200 gram). Included in Foreign Material	N/A
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable Material and all vegetable matter other than Field Pea seed material as outlined below	3
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Dead (count per 200 gram). Whole or substantially whole (more than half) including bodies	1
Field Insects Maximum Dead (count per 200 gram). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole.	15
Grasshoppers & Locusts Maximum Dead or Alive (count per 200 gram). Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces aligned end on end	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-10.1.3 Peas – Field No.1 Minimum Export Standard Machine Dressed

(Grain Trade Australia) PEAS – FIELD STANDARDS 2025/2026	
Commodity: Peas - Field Grade: No.1 Minimum Export Standard Farmer Dressed	Season: 2025/2026 Standard Reference No.: CSP- 10.1.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The peas shall be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Purity (% by weight) Minimum. Whole Field Peas, Defective Field Peas and Field Pea seed coats.	99
Moisture (%) Maximum	14
+ NOTE: For the No.1 grade, not less than 70% by weight of the entire sample shall be prime peas, that is, field peas of a size that will not pass through a 6.35mm round hole screen.	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Field Peas not of the specified variety. Field Peas that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Field Peas, whether broken or unbroken, loose seed coat and all Field Pea seed material falling through the 3.75m slotted screen. Includes Poor Colour.	4% cont + 5% bulk +
of which, Poor Colour (% by weight) Maximum. Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	1
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Is included in Total Defective
+ Note: Screen only used for assessment of No.1 grade	
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8 (including falling below the 3.75mm screen where required to be used).	0.6
Clover Burr Maximum (count per 200 gram). Included in Foreign Material	N/A
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable material and all vegetable matter other than Field Pea seed material as outlined below	1
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.1
Soil Maximum (% by weight)	
Snails Maximum Dead or Alive (count per 200 gram). Whole or substantially whole (more than half) including bodies	Nil
Field Insects Maximum Dead (count per 200 gram). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole.	1
Grasshoppers & Locusts Maximum Dead (count per 200 gram). Whole or parts thereof.	Nil
Ryegrass Ergot Maximum (length in cm). Pieces aligned end on end	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-10.2.1 Peas – Field No.2 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) PEAS STANDARDS 2025/2026	
Commodity: Peas - Field	Season: 2025/2026
Grade: No.2 Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP- 10.2.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The peas shall be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Purity (% by weight) Minimum. Whole Field Peas, Defective Field Peas and Field Pea seed coats.	97
Moisture (%) Maximum	14
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Field Peas not of the specified variety. Field Peas that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Field Peas, whether broken or unbroken, loose seed coat. Includes Poor Colour.	7
of which, Poor Colour (% by weight) Maximum. Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	N/A
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective # Is included in Total Defective	
FOREIGN SEED CONTAMINANTS Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8.	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable Material and all vegetable matter other than Field Pea seed material as outlined below	3
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Dead or Alive (count per 200 gram). Whole or substantially whole (more than half) including bodies	1
Field Insects Maximum Dead or Alive (count per 200 gram). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole.	15
Grasshoppers & Locusts Maximum Dead or Alive (count per 200 gram). Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces aligned end on end	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-10.2.2 Peas – Field No.2 Minimum Export Standard Farmer Dressed

(Grain Trade Australia) PEAS STANDARDS 2025/2026	
Commodity: Peas - Field	Season: 2025/2026
Grade: No.2 Minimum Export Standard Farmer Dressed	Standard Reference No.: CSP-10.2.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The peas shall be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Purity (% by weight) Minimum. Whole Field Peas, Defective Field Peas and Field Pea seed coats.	97
Moisture (%) Maximum	14
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Field Peas not of the specified variety. Field Peas that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Field Peas, whether broken or unbroken, loose seed coat. Includes Poor Colour.	9% Cont. 11% Bulk
of which, Poor Colour (% by weight) Maximum. Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	N/A
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective # Is included in Total Defective	
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8.	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable Material and all vegetable matter other than Field Pea seed material as outlined below	3
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Dead (count per 200 gram). Whole or substantially whole (more than half) including bodies	1
Field Insects Maximum Dead (count per 200 gram). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole.	15
Grasshoppers & Locusts Maximum Dead or Alive (count per 200 gram). Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces aligned end on end	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-10.2.3 Peas – Field No.2 Minimum Export Standard Machine Dressed

(Grain Trade Australia) PEAS STANDARDS 2025/2026	
Commodity: Field Peas	Season: 2025/2026
Grade: No.2 Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-10.2.3
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The peas shall be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Purity (% by weight) Minimum. Whole Field Peas, Defective Field Peas and Field Pea seed coats.	99
Moisture (%) Maximum	14
DEFECTIVE GRAINS -Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Field Peas not of the specified variety. Field Peas that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Field Peas, whether broken or unbroken, loose seed coat. Includes Poor Colour.	2% Cont. 5% Bulk
of which, Poor Colour (% by weight) Maximum. Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	N/A
Severely Damaged Maximum (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective # Is included in Total Defective	
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8.	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable Material and all vegetable matter other than Field Pea seed material as outlined below	1
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.1
Soil Maximum (% by weight)	0.1
Snails Maximum Dead or Alive (count per 200 gram). Whole or substantially whole (more than half) including bodies	Nil
Field Insects Maximum Dead (count per 200 gram). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole.	1
Grasshoppers & Locusts Maximum Dead (count per 200 gram). Whole or parts thereof.	1
Ryegrass Ergot Maximum (length in cm). Pieces aligned end on end	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-10.3.1 Peas - Field No.3 Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) PEAS STANDARDS 2025/2026	
Commodity: Peas - Field Grade: No.3 Minimum Receival Standard Farmer Dressed	Season: 2025/2026 Standard Reference No.: CSP-10.3.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - The peas shall be hard and well filled.	
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
Purity (% by weight) Minimum. Whole Field Peas, Defective Field Peas and Field Pea seed coats.	97
Moisture (%) Maximum	14
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Field Peas not of the specified variety. Field Peas that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Field Peas, whether broken or unbroken, loose seed coat. Includes Poor Colour.	25
of which, Poor Colour (% by weight) Maximum. Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	N/A
of which, Insect Damaged (% by weight) Maximum.	10
Severely Damaged Maximum (grains per 200 gram). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective	# Is included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8.	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material (% by weight) Maximum. Includes Unmillable Material and all vegetable matter other than Field Pea seed material as outlined below	3
Unmillable Material (% by weight) Maximum. Includes soil, stones, sclerotes and other non-vegetable material.	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum Dead or Alive (count per 200 gram). Whole or substantially whole (more than half) including bodies	1
Field Insects Maximum Dead or alive (count per 200 gram). Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole.	15
Grasshoppers & Locusts Maximum Dead or Alive (count per 200 gram). Whole or parts thereof.	2
Ryegrass Ergot Maximum (length in cm). Pieces aligned end on end	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-10.4 Peas – Yellow Split Minimum Export Standard Machine Dressed

(Grain Trade Australia) PEAS STANDARDS 2025/2026	
Commodity: Peas - Field	Season: 2025/2026
Grade: Yellow Split Minimum Export Standard Machine Dressed	Standard Reference No.: CSP-10.4
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL QUALITY PARAMETERS - Yellow Split Peas having a clean and bright appearance.	
Moisture (%) Maximum	14
Purity Minimum (% by weight) Whole peas, split peas, broken & kibbled, Defective Yellow Split Peas, Caps but excludes detached seed coats.	99.5
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight, 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Caps, Whole Field Peas with seed coat intact, discoloured yellow split peas and Whole Field peas with a missing seed coat. Includes Severely Damaged.	3
of which, Caps, Whole Field Peas with seed coat intact and discoloured yellow split peas (% by weight) Caps refer to seed coats adhering to split or broken seed.	3
of which, Whole Field Peas with a missing seed coat Maximum (% by weight). Whole Field peas with a missing seed coat	2
Detached Seed Coat Maximum (% by weight) Seed coats that are not attached to the kernels.	0.1
Broken & Kibbled Maximum (% by weight) Split pea seed material falling through the 3.97mm round hole screen	4
Poor Colour Maximum (% by weight) Yellow Split Peas whose kernels are distinctly off colour from the characteristic colour of the predominating class. Includes dark green and tinged green.	1 Dark Green 3 Tinged Green
Severely Damaged Maximum (& by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Maximum (total count per 200 gram sample; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram sample. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram sample.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Maximum (% by weight). Foreign seeds not specified in Types 1-8 falling below screen	0.6
OTHER CONTAMINANTS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Includes Unmillable material and all vegetable matter other than Yellow Split Pea material as outlined below.	0.5
Unmillable Material Maximum (% by weight) Includes soil, stones, sclerotes and other non- vegetable material. Not more than one (1) superficially obvious stone per kg.	0.1
Soil Maximum (% by weight).	
Snails Maximum Live or Dead. Whole or substantially whole (more than half) including bodies per 200 gram sample.	Nil
Field Insects Maximum Live or Dead. Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil (dead only), Sitona Weevil, Rutherglen bugs, ladybirds and wood bugs. Whole only.	Nil
Grasshoppers & Locusts Maximum Live or Dead per 200 gram sample. Whole or parts thereof.	
Ryegrass Ergot Maximum (length in cm). Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-11.1 Pigeon Peas - Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) PIGEON PEAS STANDARDS 2025/2026	
Commodity: Pigeon Peas	Season: 2025/2026
Grade: Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP 11.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS – Pigeon Peas shall be sound, dry, fresh, and light to medium brown in colour.	
Purity Minimum (% by weight). Whole Pigeon Peas, Defective Pigeon Peas and Pigeon Pea seed coats.	97
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Pigeon Peas not of the specified variety. Pigeon Peas that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Pigeon Peas, whether broken or unbroken, and loose seed coat.	5
Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	1
Severely Damaged Max (grains per 200g). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS Max (total count per 200 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynth, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8.	0.6
OTHER CONTAMINANTS Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Pigeon Pea seed material as outlined below.	3
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum. Dead or Alive. Whole or substantially whole (more than half) including bodies per 200g sample.	1
Field Insects. Maximum Dead or Alive per 200 gram sample.	15
Grasshoppers & Locusts Maximum Dead or Alive per 200 gram sample	2
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-11.2 Pigeon Peas - Minimum Export Standard Farmer Dressed

(Grain Trade Australia) PIGEON PEAS STANDARDS 2025/2026	
Commodity: Pigeon Peas	Season: 2025/2026
Grade: Minimum Export Standard Machine Dressed	Standard Reference No.: CSP 11.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS – Pigeon Peas shall be sound, dry, fresh and light to medium brown in colour.	
Purity Minimum (% by weight). Whole Pigeon Peas, Defective Pigeon Peas and Pigeon Pea seed coats.	99
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Pigeon Peas not of the specified variety. Pigeon Peas that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Pigeon Peas, whether broken or unbroken, and loose seed coat.	2
Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	1
Severely Damaged Max (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Commercial), Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Common), Oats (Sand), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum (Forage/Grain), Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Pigeon Pea seed material as outlined below.	1
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.1
Soil Maximum (% by weight)	
Snails Maximum. Dead or Alive. Whole or substantially whole (more than half) including bodies per 200g sample.	Nil
Field Insects. Maximum Dead per 200 gram sample.	1
Grasshoppers & Locusts Maximum Dead per 200 gram sample	
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

CSP-12.1 Vetch - Minimum Receival Standard Farmer Dressed

(Grain Trade Australia) VETCH STANDARDS 2025/2026	
Commodity: Vetch	Season: 2025/2026
Grade: Minimum Receival Standard Farmer Dressed	Standard Reference No.: CSP 12.1
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS – Vetch shall be sound, dry, fresh and colour typical of the variety of the season.	
Purity Minimum (% by weight). Whole Vetch, Defective Vetch and Vetch seed coats.	97
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Vetch not of the specified variety. Vetch kernels that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Vetch, whether broken or unbroken and loose seed coat. Vetch where whole or part of the seed coat only is damaged, is included as sound Vetch.	5
Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	1
Severely Damaged Max (grains per 200g). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 ^
^ Not included in Total Defective # Included in Total Defective	
FOREIGN SEED CONTAMINANTS Max (total count per 200 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Oats (Common), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8.	0.6
OTHER CONTAMINANTS Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Vetch seed material as outlined below.	3
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.5
Soil Maximum (% by weight)	0.3
Snails Maximum. Dead or Alive. Whole or substantially whole (more than half) including bodies per 200g sample.	1
Field Insects. Maximum Dead or Alive per 200 gram sample.	15
Grasshoppers & Locusts Maximum Dead or Alive per 200 gram sample	2
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 200 gram sample.	2
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	1
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil

CSP-12.2 Vetch - Minimum Export Standard Machine Dressed

(Grain Trade Australia) VETCH STANDARDS 2025/2026	
Commodity: Vetch	Season: 2025/2026
Grade: Minimum Export Standard Machine Dressed	Standard Reference No.: CSP 12.2
Sampling Guidelines	
A minimum of 3 probes (litres) must be taken from every load tendered for delivery, plus an additional (probe) litre for every 10 tonnes in the load.	
PHYSICAL CHARACTERISTICS - Vetch shall be sound, dry, fresh and be colour typical for the variety of the season.	
Purity Minimum (% by weight). Whole Vetch, Defective Vetch and Vetch seed coats.	99.5
Moisture (%) Maximum	14
Approved Varieties. Approved varieties as listed in these Standards or on the Grain Trade Australia website in the Masterlist.	
DEFECTIVE GRAINS - Maximum (% by weight in 200 gram sample; unless otherwise stated)	
Defective Maximum (% by weight). Vetch not of the specified variety. Vetch kernels that are broken, chipped, diseased, frost damaged, insect damaged, sappy, shrivelled, split, sprouted, weather damaged, wrinkled. Includes pods that contain Vetch, whether broken or unbroken and loose seed coat. Vetch where whole or part of the seed coat only is damaged, is included as sound Vetch.	5
Poor Colour Maximum (% by weight). Seed coat or kernel that is distinctly off colour from the characteristic colour of the predominating class.	1
Severely Damaged Max (% by weight). Mould, Burnt & Heat Damaged or Other Serious Visual Defects.	1 #
^ Not included in Total Defective	# Included in Total Defective
FOREIGN SEED CONTAMINANTS - Max (total count per 200 gram; unless otherwise stated) - Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per 200 gram. Except Type 1 in which the maximum applies on an individual seed basis per 200 gram.	
Type 1 Max* (Individual seeds per 200 gram). Colocynthis, Double Gee/Spiny Emex/Three Cornered Jack, Jute, New Zealand Spinach, Parthenium Weed (QLD only. NIL tolerance in NSW/VIC/SA), Poppy (Field), Poppy (Horned), Poppy (Long Head), Poppy (Mexican), Poppy (Wild)	4*
Type 2 Max (entire load). Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Opium Poppy, Peanut Seeds & Pods, Ragweed, Rattlepods, Starburr, St John's Wort	Nil
Type 3a Max (seeds in total per 200 gram). Bathurst Burr, Branched Broomrape, Bulls Head/Caltrop/Cats Heads, Cape Tulip, Cotton Seed, Dodder, Noogoora Burr, Thornapple	1
Type 3b Max (seeds in total per 200 gram). Vetch (Tare)	2
Type 3c Max (seeds in total per 200 gram). Heliotrope (Blue), Heliotrope (Common). Pods must be opened, and seeds counted	4
Type 4 Max (seeds in total per 200 gram). Bindweed (Field), Cutleaf Mignonette, Darnel (Drake Seed), Hexham Scent/Melilot (King Island) (only acceptable if no tainting odour is present), Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	10
Type 5 Max (seeds in total per 200 gram). Knapweed (Creeping/Russian), Pattersons Curse/Salvation Jane, Sesbania Pea	20
Type 6 Max (seeds in total per 200 gram). Clover (Pods), Columbus Grass, Johnson Grass, Lucerne (Pods), Marshmallow (Pods), Medic (Pods), Muskweed (Pods), Saffron Thistle, Sow Thistle Heads, Trefoil (Pods), Wild Radish (Pods)	5
Type 7a Max (seeds in total per 200 gram). Adzuki Beans, Broad Beans, Chickpeas (Desi/Kabuli), Corn (Maize), Cowpea, Faba Beans, Lentils (Green/Red), Lupin (Albus/Angustifolius), Mungbeans, Onion Weed pods, Peas (Field), Pigeon Peas, Soybean. Any other seeds or pods greater than 5mm diameter	10
Type 7b Max (seeds in total per 200 gram). Barley, Bindweed (Australian), Bindweed (Black), Brome Grass, Carrot Weed, Durum, Oats (Black/Wild - individual seeds counted in clusters), Oats (Sand), Oats (Common), Rice, Rye (Cereal), Ryegrass on Stalk, Sorghum, Triticale, Turnip Weed Pods, Wheat. Any other seed not specified in TYPE 1 to TYPE 7(a), TYPE 7(c), TYPE 8 or SFS	10
Type 7c Max (seeds in total per 200 gram). Safflower, Sunflower	1
Type 8 Max (seeds in total per 200 gram). Bellvine	100
Small Foreign Seeds Max (% by weight). Foreign seeds not specified in Types 1-8.	0.6
OTHER CONTAMINANTS - Max (% by weight in 200 gram sample; unless otherwise stated)	
Foreign Material Maximum (% by weight). Unmillable material and all vegetable matter other than Vetch seed material as outlined below.	0.5
Unmillable Material Maximum (% by weight). Soil, stones, sclerotes and non-vegetable matter	0.1
Soil Maximum (% by weight)	
Snails Maximum. Dead or Alive. Whole or substantially whole (more than half) including bodies per 200g sample.	Nil
Field Insects. Maximum Dead per 200 gram sample.	1
Grasshoppers & Locusts Maximum Dead per 200 gram sample	
Ryegrass Ergot Maximum (length in cm) Pieces laid end to end per 200 gram sample.	Nil
Gumnuts Maximum (Count per 2kg). Whole or pieces of gumnut of any size and maturity level.	Nil
Objectionable Material Maximum (entire load). Includes animal excreta, rodents, crushed insect bodies or parts that adhere to the grain, live stored grain insect pests, live pea weevil, pickling compounds, tainting agents, odours or any other commercially unacceptable contaminant, smell, or taste.	Nil
# When not in conflict with Storage and Handling Agreement or Marketing Contract. Please note that this Export Standard is in addition to the specific quarantine requirements of particular countries, as governed by Plant Exports Branch.	

SECTION 4 VARIETY CLASSIFICATION

At present, there are no varietal restrictions regarding any pulses listed in these Standards.

When applying these Standards, industry should note:

- For some pulse commodities, Standards may apply by variety. In that instance, a separate segregation may be required for each individual variety. Standards may or may not include a minimum varietal purity requirement.
- For most pulse commodities, there are no varietal restrictions, and all varieties may be received into a grade and segregation.
- Some markets required particular sizes or shapes of particular pulses. Industry advice on the GTA website of the size and shape of certain pulses (i.e., red lentils) is available for guidance.

SECTION 5 METHODS & PROCEDURES

5.1 Introduction

The following section details methods and procedures to be used for the assessment of various quality parameters as outlined in this Manual.

The methods outlined are Reference Methods, Field Assessment Methods or suggested method of analysis for various quality parameters. Field Assessment Methods are included as a guide to industry where Reference Methods may not be able to be implemented. Note that Field Assessment Methods must equate to the Reference Method for the applicable test method.

In all instances of disputes, test results produced by trade-certified equipment take precedence over non-trade certified equipment and methods. Where the dispute involves only non-trade-certified equipment or test methods, the reference method takes precedence over the field assessment method.

Depending on the test to be conducted, variations may exist due to equipment used.

Procedures outlined are a guide for industry. Industry is free to develop their own Operational Procedures for each test and activity based on their own circumstances. At all times industry use of apparatus outlined in this Standard must comply with the manufacturers' recommendations for occupational health and safety and training.

5.2 Sampling

5.2.1 Definitions

This is the standard procedure used to draw a sample of the commodity from a bulk unit tendered for delivery to enable tests to be conducted on the commodity for the purposes of determining its quality.

- A primary sample is an individual probed sample taken from the lot presented for sampling.
- A composite sample is the combined primary samples taken from the lot to be sampled and is representative of the entire lot.
- A sub sample is the sample taken from the mixed composite sample for the purposes of conducting quality tests and is representative of the entire lot.

5.2.2 Scope

Pulses are traded on the basis of quality tests conducted on lots of the pulse presented for sale or delivery to end users. Obtaining representative samples is critical to ensuring test results reflect the true quality of these lots.

This procedure is applicable to all pulses.

5.2.3 Apparatus

- Manual sampling probe (double tube compartment probe, one inside the other, equipped with spiralled ports that open sequentially from bottom to top).
- Vacuum or pneumatic probe (an alternative to the manual sampling probe and consisting of a handheld or remotely controlled probe which retrieves grain through the use of a vacuum or other air movement system).
- Any other sampling device capable of obtaining a representative sample of pulses.
- Mixing bucket (including other associated equipment such as mini-auger suitable for mixing sample, optional).
- Sample dividing apparatus (optional).

5.2.4 Reagents

Not Applicable.

5.2.5 Procedure

Sample Collection guidelines for collecting a representative sample.

- The surface of the grain should be fully exposed prior to sampling to allow for effective visual inspection. At this point, the load should be scanned for any defects or contaminants.
- The probe to be used should be of a sufficient length in order to obtain a sample from as close as possible to the bottom of truck.
- A primary sample must be drawn for assessment by thrusting the sampling probe as vertically and as deep as possible into the load.
- At least one probe must be taken from the front, middle and rear of each bulk unit.
- If more than one unit is delivered, samples must be drawn from each bulk unit as described above.
- If the bulk units are of visibly different quality, or if required at the Receiving Agents discretion, different samples and grade classification may be undertaken for each separate bulk unit.
- If the declared varietal composition or paddock where the grain was grown is different for each unit tendered for delivery, or more than one variety is commingled in each delivery unit, then a separate assessment of each unit must be conducted.

- Each primary (probed) sample must consist of at least one litre of grain.
- A composite sample from each load tendered for delivery shall consist of the following minimum quantities and number of probes:

Load Size	Sample Size (minimum)
10 tonnes or less	3 litres
Over 10 tonnes up to 20 tonnes	4 litres
Over 20 tonnes up to 30 tonnes	5 litres
Over 30 tonnes up to 40 tonnes	6 litres
Over 40 tonnes up to 50 tonnes	7 litres
Over 50 tonnes up to 60 tonnes	8 litres
Over 60 tonnes up to 70 tonnes	9 litres
Over 70 tonnes up to 80 tonnes	10 litres

Note – in the above table the sample size reflects the number of probe samples. For example, 4 litres equates to 4 probe samples.

Sample Mixing

- The primary samples in each probe must be collected together and thoroughly mixed in a suitable container using a mechanical device where appropriate, to form the composite sample.
- Where large samples are obtained in this manner or where high levels of Foreign Material are present in the sample, it is recommended that a suitable mechanical device is used for mixing and sub-dividing the sample.
- Sub samples should be drawn from the composite sample either by hand or through the use of a suitable sample dividing apparatus.

Sample Analysis

- The sub-sample obtained for assessment should be a representative 200 gram, 400 gram or 600 gram sample depending on the grain type as per the following list:

200 gram Sample	400 gram Sample
Adzuki Beans	Chickpeas – Kabuli Type No. 1 Grade Large*
Caloona / Poona Cowpeas	Faba Beans – whole and split
Chickpeas – Desi Type	Lupins – Albus
Chickpeas – Split Chana Dhal	
Chickpeas – Kabuli Type No. 1 Grade Small*	
Fenugreek	
Lentils – Whole Green	
Lentils – Whole Red	
Lentils – Split Red	
Lupins – Angustifolius	
Mung Beans	600 gram Sample
Peas – Field	Broad Beans – whole and split
Peas – Yellow Split	
Pigeon Peas	
Vetch	

* Varieties are classified as Small or Large based on size

- The sub sample should then be analysed for all of the quality parameters specified in the applicable pulse Standards or in the Receiving Agent's agreement with the buyer concerned if different from these Standards.
- Results should be entered on the Receiving Agents sample receipt.

5.2.6 References

Sampling of Wheat and other Grains - AACC Method 64-70A

5.3 Moisture Assessment - NIR

5.3.1 Definitions

This describes the NIR method for determination of moisture in all pulses.

5.3.2 Scope

This procedure is applicable to all pulses.

5.3.3 Reagents

Not applicable.

5.3.4 Apparatus

NIR instrument approved for use for pulses with an appropriate calibration.

5.3.5 Method

Sample to be “as is”.

Individual manufacturer instructions and procedures should be followed for operation and maintenance of NIR instruments used to determine grain moisture.

Report result to the nearest 0.1%.

5.3.6 References

- Grain industry Code of Practice

5.4 Defective Grains Assessment – Reference Method

5.4.1 Definitions

This describes the method of assessment of deliveries of pulses for the various types of defective grains described in these pulse Standards. Refer also to the de-hulling procedure.

5.4.2 Scope

This method is applicable for all deliveries of pulses.

5.4.3 Apparatus

Refer to the screen to be used for the assessment of defective grains in each applicable commodity Standard, noting there are no reference screen specifications for pulse screens.

Visual Recognition Standards, with the following photographic standards being recognised by GTA:

- Visual Recognition Standards Guide for Grain Commodity Sampling and Assessment – Issued August 2025, GTA
- Australian Pulse Varieties 2024, Grains Australia

5.4.4 Reagents

Not applicable

5.4.5 Method

General

- Sample to be “as is”.
- Note that variations to the assessment of Defective grains as outlined in the following procedure may apply depending on the pulse commodity.
- Where required, depending on the Defect and pulse commodity, the seed coat must be removed to examine the kernel for the presence of the Defect. This process is referred to as De-hulling. Refer to the De-Hulling method.
- Each type of defect can be assessed with the assistance of the GTA Approved photographic standards (Visual Recognition Standards Guide) or Approved objective measurement instruments where appropriate.
- Instruments of magnification may be used to aid the assessment of pulses for Defective Grains.
- The time taken to assess the sample for defects is unlimited.
- Samples should be assessed under conditions of good lighting.
- Obtain a 200 gram, 400 gram or 600 gram sample to the nearest 0.1 gram, as applicable based on the pulse commodity.
- Examine material in the entire sample (including all trays) for the presence of Defects for which a nil tolerance applies. For nil tolerance defects, the tolerance (e.g., rejection of the load) can apply if the defect is detected at any stage of the delivery or testing process, including in the truckload before sampling, in the probe sample, in the entire subsample or during discharge into the receival hopper after assessment.

- The defective seed category includes all the defective seeds defined in the comments / variations section of each Standard.
- Where a pulse commodity has a defect listed in a separate category in the Standards, these must be counted separately. Where a Total Defective category exists and the separate category specifies inclusion in the Total Defective category, they must be counted in that Total Defective category after counting for that separate sub-category.
- An individual seed or kernel may only have one defect, being the defect type with the tightest tolerance in the Standard.
- Both sides of the grain should be inspected to determine firstly whether a Defect is present and secondly to determine if its presence is as per the Definitions to classify as Defective.
- Where low but acceptable levels of defects such as Fungal Affected lesions (e.g., Ascochyta) are found on the seed coat, it may be desirable for seed coats to be removed to determine the presence on the kernel.
- If a separate tolerance for particular defects such as Fungal Affected (e.g., Ascochyta) or Poor Colour is listed in the Standard, examine all the relevant trays or the entire 200 gram, 400 gram or 600 gram sample for the presence of the defect as defined in the Standard.
- For Severely Damaged, the entire 200 gram, 400 gram or 600 gram sample is to be assessed for most pulse commodities:
 - When assessing as per the Receival Standard, the number of grains in the entire sample with this defect are counted. For most pulse commodities, the level found is not included in Total Defectives.
 - When assessing as per the Export Standard, the weight of grains as a percentage of the entire sample is to be determined. For most pulse commodities, the level found is included in Total Defectives.

Varietal Restriction

- For a list of approved varieties by commodity, refer to the Varietal Masterlist on the Grain Trade Australia website at www.graintrade.org.au.
- For those commodities where a tolerance applies, examine material in the entire sample (including all trays) for the presence of varieties not of the declared variety.

Contrasting Colour

- For those commodities where a tolerance applies, examine material in the entire sample (including all trays) for the presence of Contrasting Colour. Refer to the reference Variety Colour Charts or GTA Visual Recognition Standards Guide to assist.
- Where low but acceptable levels are found on the seed coat, it may be desirable for seed coats to be removed to determine the presence on the kernel.

Where a Size Limit Applies i.e., a Screen is specified

- Place the entire 200 gram, 400 gram or 600 gram sample on the top of the appropriate screen(s). The screen size for those commodities where a screen is required for assessment is listed in the applicable Standard.
- For some commodities a different screen size is to be used based on the variety.

- Screen the sample using the Sieving Method procedure.
- Any pulse material (defined as the entire seed, pieces of kernel with/without the seed coat attached and seed coats) of the pulse type being assessed that fall through the screen, including whole grains, are classified as defective.
- Examine the top tray (or middle tray if applicable) for the various Defect types, pick out and weigh each Defect subcategory as listed in each Standard.

Where a Size Limit Does Not Apply i.e., a Screen is not specified

- Hand-pick any Defective pulses from the entire 200 gram, 400 gram or 600 gram sample.
- Calculate results as follows:
 - For each individual defect with a % by weight tolerance, pick out the defective grains from the entire sample and weigh. Divide the weight by two for a 200 gram sample, by four for a 400 gram or by six for a 600 gram sample to calculate the % of the defective grains. This should generally be done before the assessment of Total Defects in the sample.
 - The percentage of Total Defects is calculated based on the weight of the individual defects combined or Total of all defects as listed where applicable in each Standard in the entire 200 gram, 400 gram or 600 gram sample.
 - For those defects with a maximum number in the entire sample, pick out those defective grains and count them.
 - Report all applicable results to the nearest 0.1% or nearest number per sample size (200 gram, 400 gram or 600 gram) whichever is applicable.

5.4.6 References

Visual Recognition Standards Guide for Grain Commodity Sampling and Assessment – Issued August 2025, GTA.

Australian Pulse Varieties 2024, Grains Australia.

5.5 Contaminants & Purity Assessment – Reference Method

5.5.1 Definitions

This describes the method of assessment of deliveries of pulses for the various types of Contaminants described in the relevant pulse Standard. In general, these are called Foreign Material.

Purity is defined as all pulse seed material of the pulse being assessed. This includes whole pulse seed, defective pulse seed and pulse Seed Coats. Purity is the opposite of Foreign Material.

The various contaminant types and their assessment methods as described in this method are as follows:

% by weight in entire sample	Count per entire sample	Count per entire load	Length in cm per entire sample	Count per 2kg
Small Foreign Seeds	All Weed Seeds except Type 2	Chemicals not approved for pulses or in excess of the MRL	Ryegrass Ergot	Gumnuts
Foreign Material	Snails	Objectionable Material		
Unmillable Material	Field Insects	Type 2 Weed Seeds		
Soil	Grasshoppers, Locusts			
Purity*				

* Not a contaminant. Purity is the opposite of Foreign Material

5.5.2 Scope

This method is applicable for all deliveries of pulses.

5.5.3 Apparatus

Refer to the screen to be used for the assessment of defective grains in each applicable pulse Standard.

Analytical balance accurate to at least 0.01g.

Visual Recognition Standards with the following photographic standards being recognised by GTA:

- Visual Recognition Standards Guide for Grain Commodity Sampling and Assessment – Issued August 2025, GTA.
- Seed Impurities of Grain Identification Guide, 3rd Edition, GTA.
- Insects of Stored Grain, A Pocket Reference, 2nd Edition, CSIRO

A tray or mechanism capable of holding the relevant sample size.

Colander or Mesh Screen (optional).

Ruler.

5.5.4 Reagents

Not applicable.

5.5.5 Method

General

- Sample to be “as is”.
- Note that variations to the assessment of Contaminants and Purity as outlined in the following procedure may apply depending on the pulse commodity.
- Note that instruments of magnification may be used to aid the assessment of pulses for Contaminants and Purity.
- Samples should be assessed under conditions of good lighting.
- To assist in the separation of the pulse material (Purity) from Contaminants (all other material) in the sample, various screens may be used. After sieving, the sample may still need to be hand-picked to separate the various fractions of contaminants. Refer to the Sieving Procedure.
- The time taken to assess the sample for Contaminants and Purity is unlimited.

Procedure

- Obtain a 200 gram, 400 gram or 600 gram sample to the nearest 0.1 gram, as applicable based on the pulse commodity.
- Examine material in the entire sample (including all trays where a sieve is required to be used) for the presence of Contaminants for which a nil tolerance applies. For nil tolerance Contaminants, the tolerance (e.g., rejection of the load) can apply if the Contaminant is detected at any stage of the delivery or testing process, including in the truckload before sampling, in the probe sample, in the entire subsample or during discharge into the receival hopper after assessment.
- Examine the entire sample and pick out any Contaminants. Where individual tolerances exist for categories within Foreign Material, these must be separated from the sample into their individual constituents. It may be easier to weigh each category separately to determine the level prior to combining all categories to determine the total Foreign Material percentage by weight.
- From the total Foreign Material, extract the Unmillable Material (or alternatively conduct this assessment prior to determining total Foreign Material). Using a device such as a colander, sieve this Unmillable Material to remove the soil. It may be necessary to continue to hand-pick soil from the sample to remove it all.
- Examine all the trays or the entire 200 gram, 400 gram or 600 gram sample for the presence of nominated Foreign Seeds. Any Foreign Seed pods must be opened and the seeds counted except where pods have a specified tolerance. Most commodities have common tolerances for seeds in the entire 200 gram, 400 gram or 600 gram sample. Note that for Type 1 weed seeds, tolerances apply to individual seeds whereas for all other Types listed, tolerances are the total of all seeds in each Type. Refer to the relevant pulse Standard for tolerances to apply.
- From the Foreign Seeds category, separate out any Small Foreign Seeds (generally collected from the bottom catch pan) where a sieve is used, or from the total sample if a sieve is not used, and weigh them. Again, a colander or mesh may assist in this task, but hand-picking may still be required.
- For all pulses, where Field Peas are present in the sample, it is recommended these be broken open to determine the presence of live Pea Weevil. Where required additional Field Peas should be collected from the probe sample or from the surface of the truck. When assessing the commodity Field Peas, a representative number should be selected from the Field Pea sample and broken open to determine the presence of live Pea Weevil.

- For Ryegrass Ergot, generally collected in the bottom catch pan where a sieve is used, align all pieces end on end and measure the length in cm. This material is included in the total Foreign Material.
- If a Gumnut is found in the initial 200 gram, 400 gram or 600 gram sample, then a further sample should be taken to reach a weight of 2kg. Count the total number of whole or pieces of Gumnuts as per the relevant Standard.
- Where depicted, other Contaminants should be assessed using the GTA Approved photographic standards. Where reference material is not available, other Contaminants should be assessed by reference to the Definitions of those parameters in these Standards.
- For assessment of Pickling Compounds or Artificial Colour, Chemicals not Approved for pulses or Chemicals in Excess of the MRL, it is recommended that all deliveries are accompanied by a signed declaration referring to its chemical status. Where the receiving agent believes that the visual appearance and/or odour of grain suggests that it has been treated with a non-approved chemical, it is recommended the grain is not received until the representative “as received” sample has been tested by an approved independent laboratory and the presence or absence of non-approved chemicals ascertained.

Reporting Results

For most Contaminants, divide the weight of each Contaminant category and total weight of all Contaminant categories combined by two for a 200 gram sample, by four for a 400 gram or by six for a 600 gram sample to calculate the % of each category and Foreign Material level combined.

For Purity, divide the total weight of all the applicable pulse being assessed material combined by two for a 200 gram sample, by four for a 400 gram or by six for a 600 gram sample to calculate the %. Alternatively, Purity is the opposite of Foreign Material (i.e., $100 - \text{FM}\%$).

For Ryegrass Ergot, record to the nearest 0.1cm.

For recording, report results as follows:

Count per entire sample – nearest whole number
Length in cm per half litre – nearest 0.1cm
Percentage by weight in entire sample – nearest 0.1%

5.5.6 References

Visual Recognition Standards Guide for Grain Commodity Sampling and Assessment – Issued August 2025, GTA.

Seed Impurities of Grain Identification Guide, 3rd Edition, GTA

Insects of Stored Grain, A Pocket Reference, 2nd Edition, CSIRO

Ute Guide Series, GRDC.

5.6 Sieving Method

5.6.1 Definition

This procedure is to be used for the assessment of Defective grains where for the pulse in question being assessed, all pulse seed material that falls below the screen is included in the definition for Defective.

See also the definition for Seed Material.

5.6.2 Scope

This procedure is applicable to all pulses where all pulse seed material falling below the screen is included in the definition for Defective.

5.6.3 Apparatus

Shaking Device

Refer to the screen to be used for the assessment of defective grains in each applicable commodity Standard, noting there are no reference screen specifications for pulse screens.

Analytical balance accurate to at least 0.01g.

5.6.4 Reagents

Not applicable.

5.6.5 Procedure

- Either an automatic shaking machine or a manual shaking process may be used. In either scenario, the minimum number of shakes is ten.
- Note that during the process of shaking some commodities such as split product, care must be taken to not increase the level of defectives such as broken and splits (kibbled).

Automatic Shaking Device

If an automatic shaking machine is used, it must correlate to the Manual Reference Method below.

Manual Shaking Reference Method

The Manual Reference Method is as follows:

- a. A 200 gram, 400 gram or 600 gram sample of grain shall be weighed on an appropriate balance that is accurate under the conditions of use to plus or minus 0.01 gram.
- b. All screen surfaces (top, middle and/or bottom) shall be clean, smooth, dry and free of grain residue.
- c. The sieving process shall occur on a flat and smooth screen movement table. Marks at each end of the table that indicate a screen movement of thirty centimetres may assist.
- d. With the screen resting against one of the marks, the 200 gram, 400 gram or 600 gram sample of grain is to be placed in one movement centrally onto the surface of the screen. No additional movement or spreading of the sample over the screen surface is to occur.
- e. Where applicable, with the slots facing away/towards the operator (top, middle and/or bottom), the screen shall be evenly moved ten times in a to and fro motion, that is, ten aways and ten returns in the direction of the slots and with the screen being moved on the surface of the screen

movement table.

- f. Each of the ten to and fro movements are to take one second, so that the complete screen movement process occupies ten seconds.
- g. The front edge of the screen shall travel thirty centimetres forward and the same distance back.
- h. At the completion of the ten to and fro movements, the screen shall be gently removed from the bottom catch pan.
- i. Pulse seed material passing through the nominated screen into the catch pan must be separated into the various fractions as described by the relevant commodity Standard.
- If required, calculate the percentage pulse material below the screen as follows:

$$\text{Pulse material below the screen by wt. (\%)} = \frac{\text{Pulse material below the screen weight}}{\text{Total sample weight}} \times 100$$

Alternatively, the Pulse material below the screen may need to be added to the defective material retained above the screen to determine the Total Defective, depending on the commodity Standard.

5.6.6 References

Not applicable.

5.7 Sizing Method

5.7.1 Definition

This procedure is to be used for sizing of samples often required for Machine Dressed Export product. This is also referred to as Retention.

5.7.2 Scope

This procedure is applicable to all pulses where there is a requirement to size pulses.

5.7.3 Apparatus

Shaking Device

Refer to the screen to be used for the assessment of defective grains in each applicable commodity Standard, noting there are no reference screen specifications for pulse screens.

Refer also to the contract or relevant Standard which specifies a particular pulse commodity size (or screen size) for the pulse in question, noting there are no reference screen specifications for the pulse sizing screen that may be required to be used.

Analytical balance accurate to at least 0.01g.

5.7.4 Reagents

Not applicable.

5.7.5 Procedure

- While this procedure may not be required for all Machine Dressed Export product the following is the method to be used where required.
 - a. Sample to be sized is first to be assessed as per the relevant Standard.
 - b. Prior to sizing, all non-pulse material (i.e., Foreign Material) is to be removed from the sample. Weigh the remaining pulse material and record the result.
 - c. The remaining pulse material after removal of Foreign Material is to be placed on the relevant sized screen.
 - d. The screen is to be shaken using the procedure for assessment of defects falling through the screen except that the number of shakes is 40 for sizing i.e., forty shakes of the sieve using the method outlined in Procedure 5.6.
 - e. Weigh the pulse material passing through the sieve after completion of the 40 shakes.
 - f. Calculate the percentage by weight retained above the screen (Retention) using the total sample weight and the weight of pulse material falling through the screen:

$$\text{Pulse material above the screen by wt. (\%)} = \frac{\text{Pulse material above the screen weight}}{\text{Total sample weight}} \times 100$$

5.7.6 References

Not applicable.

5.8 De-hulling Method

5.8.1 Definition

This procedure is to be used for de-hulling the kernel prior to assessment of seed kernel defects as per the relevant pulse Standard for all commodities where kernels are to be assessed underneath the Seed Coat.

5.8.2 Scope

This procedure is applicable to all pulses where there is a requirement to determine the presence of a defect on the kernel via removal of the Seed Coat.

5.8.3 Apparatus

A suitable de-hulling device.

Visual Recognition Standards Guide for Grain Commodity Sampling and Assessment – Issued August 2025, GTA.

Analytical balance accurate to at least 0.01g.

5.8.4 Reagents

Not applicable.

5.8.5 Procedure

- While this procedure may not be required for all Farmer or Machine Dressed product the following is the method to be used where required.
- This procedure is only for the assessment of the defect on the kernel and is generally only applicable to determine Poor Colour kernel. Other defects, unless specified in the Standard, are generally not to be calculated based on this method. However, variations may exist.
- Assessment for kernel defects is to occur on an ‘as is’ dirty sample basis i.e., without the removal of Foreign Material or defects.
- A minimum of 200 grams of the ‘as is’ sample obtained following probing the load is to be de-hulled. A larger sample may be used where required.
- To assist in a more rapid assessment process, the de-hulling method should occur at the same time as the sample is being assessed for other quality parameters i.e., obtain a minimum 200 grams from the probe sample and start the de-hulling unit and a further 200 grams, 400 grams or 600 grams from the probe sample and commence assessment for all quality parameters as listed in the relevant Standard except for specific kernel defects such as Poor colour.
- Following de-hulling, weigh the pulse material that has been de-hulled i.e., all kernels, not the removed Seed Coats.
- During the de-hulling process:
 - Parts of the kernel may be broken off causing “a chalky appearance” on the kernel. Kernels are not to be assessed as defective as a result of these chalky/dusty particles. These particles and dust may be gently removed by various means such as a damp cloth.
 - Pieces of Seed Coat may be retained on the de-hulled kernel.
- Assessment for defects on the kernel is to occur on the de-hulled sample. Utilise the Visual Recognition Standards Guide where defects on the kernel may be visually defined or select material as defined in the relevant Standard.
- Weigh the pulse material defined as the relevant defect as identified following de-hulling.

- Calculate the percentage by weight of the relevant defect on the kernel following de-hulling as follows:

$$\text{Kernel defect by wt. (\%)} = \frac{\text{Kernel defect weight}}{\text{Total sample weight (200g)}} \times 100$$

- If the relevant Standard specifies, this kernel defect material may need to be included in the calculation for Total Defective.

5.8.6 References

Visual Recognition Standards Guide for Grain Commodity Sampling and Assessment – Issued August 2025, GTA.

5.9 Variety Declaration Procedure

5.9.1 Definitions

This is the recommended procedure for determining the variety of the load presented for delivery.

5.9.2 Scope

This procedure is applicable to all pulse deliveries where the variety is required to be declared or there is a limit to the admixture of more than one variety in the relevant Standard.

5.9.3 Apparatus

Not applicable.

5.9.4 Reagents

Not applicable.

5.9.5 Method

- Generally, the variety is only required to be declared under the Receival Standard at the point of delivery of the relevant pulse commodity. However, there may be a requirement to declare the variety of the pulse consignment in the Export Standard.
- For all deliveries, it is recommended the driver or deliverer of the grain declares the variety(s) in the load tendered for delivery. It is recommended that the grower sign a Declaration Form and provide this to the driver for provision to the Receival Agent. This Declaration Form should at a minimum contain the grower details and the variety(s) of the load.
- If the declared varietal composition or paddock where the grain was grown is different for each unit tendered for delivery, or more than one variety is commingled in each delivery unit, then a separate assessment of each unit must be conducted.
- While this procedure generally only applies to those commodities where the Standard specifies a limit on the admixture of more than one variety in a load, it may also be applied where there is a contractual obligation based on variety.
- Note that depending on the varietal declaration and the procedures of the Receival Agent, a sample of the load may be taken and sent to a laboratory for assessment of the variety within the sample. In this instance sample is to be “as is”.
- Report the variety as per the following procedure using the applicable code as defined by the Receival Agent.

Load is Declared as One Variety Only

- Where the load is declared as being of the one variety only, and based on the quality assessment confirming this declaration, grade the load and record the declared variety.
- Where the load is declared as being of the one variety only and based on the quality assessment more than one variety is determined to be present:
 - If there is no varietal restriction or admixture limit listed in the Standard, no further assessment is required, noting the Receival Agent may require further assessment based on their own commercial procedures.
 - If there is a varietal restriction or admixture listed in the Standard, remove and weigh the grains determined to be the additional variety.

- Record the declared variety and as relevant, the varietal admixture percentage based on the relevant calculation specified at the end of section 5.9.5.

Load is Declared as Multiple Varieties

- Where the load is declared as being of more than one variety and based on the quality assessment more than one variety is determined to be present:
 - If there is no varietal restriction or admixture limit listed in the Standard, no further assessment is required, noting the Receival Agent may require further assessment based on their own commercial procedures. Based on the quality results, Grade the load and record the variety with the greatest percentage in the load.
 - If there is a varietal restriction or admixture listed in the Standard, remove and weigh the grains determined to be the additional variety(s). Record the variety with the greatest percentage in the load and as relevant, the varietal admixture percentage based on the relevant calculation specified at the end of section 5.9.5.
- Where required, calculate the percentage by weight of the variety(s) as follows:

$$\text{Variety by wt. (\%)} = \frac{\text{Variety weight}}{\text{Total sample weight}} \times 100$$

- If the relevant Standard specifies, this varietal admixture material may need to be included in the calculation for Total Defective.

5.9.6 References

Australian Pulse Varieties 2024, Grains Australia

Declaration Form, if applicable

SECTION 6 REFERENCE MATERIALS

At the time of publishing this Manual, the following photographic Reference Material referred to in this Manual is considered by GTA to be suitable as an aid to classification of pulses.

Industry should be aware that all such material is controlled by the author of that material and appropriate copies of that material can be obtained from the author.

The method of printing, copying, storing, using or otherwise obtaining such Reference Material may impact on the appearance of its content. This may impact on the classification of pulses. Industry should note the method of publication of the material by the author and other relevant information such as version number to ensure they have the appropriate version.

Name of Material	Material Type	Author	Version Number	Applicable Dates
Defective Grains				
Visual Recognition Standards Guide for Grain Commodity Sampling and Assessment	Hardcopy booklet	GTA	n/a	Issued August 2025
Australian Pulse Varieties 2024	Soft copy	Grains Australia	2024 version	Issued 2024
Contaminants				
Grain Quality Winter Grain Crops: The Ute Guide	Hardcopy booklet	GRDC	n/a	n/a
Weeds: The Ute Guide	Hardcopy booklet	GRDC	Various editions	n/a
Insects of Stored Grain, A Pocket Reference	Hardcopy booklet	CSIRO	2 nd Edition	2007
Seed Impurities of Grain Identification Guide	Hardcopy booklet	GTA	3 rd Edition	n/a

